

INTERIOR SECOND DERIVATIVE ESTIMATES FOR SOLUTIONS TO THE LINEARIZED MONGE–AMPÈRE EQUATION

CRISTIAN E. GUTIÉRREZ AND TRUYEN NGUYEN

ABSTRACT. Let $\Omega \subset \mathbb{R}^n$ be a bounded convex domain and $\phi \in C(\overline{\Omega})$ be a convex function such that ϕ is sufficiently smooth on $\partial\Omega$ and the Monge–Ampère measure $\det D^2\phi$ is bounded away from zero and infinity in Ω . The corresponding linearized Monge–Ampère equation is

$$\text{trace}(\Phi D^2 u) = f,$$

where $\Phi := \det D^2\phi (D^2\phi)^{-1}$ is the matrix of cofactors of $D^2\phi$. We prove a conjecture in [GT] about the relationship between L^p estimates for D^2u and the closeness between $\det D^2\phi$ and one. As a consequence, we obtain interior $W^{2,p}$ estimates for solutions to such equation whenever the measure $\det D^2\phi$ is given by a continuous density and the function f belongs to $L^q(\Omega)$ for some $q > \max\{p, n\}$.

1. INTRODUCTION

L^p -estimates play a fundamental role in the theory of second-order elliptic partial differential equations, with many works devoted to the topic, see [GiT, Chapter 9] and [CC, Chapter 7]. For linear equations of the form $\text{trace}(A(x)D^2u(x)) = f(x)$ in a domain $\Omega \subset \mathbb{R}^n$ with

$$(1.1) \quad \lambda|\xi|^2 \leq \langle A(x)\xi, \xi \rangle \leq \Lambda|\xi|^2 \quad \text{for all } x \in \Omega \text{ and } \xi \in \mathbb{R}^n,$$

L^p -estimates for second derivatives of solutions were derived in the 1950's as a consequence of the celebrated Calderón and Zygmund theory of singular integrals. Precisely, if the matrix $A(x)$ is continuous in Ω , then for any domain $\Omega' \Subset \Omega$ and any $1 < p < \infty$ we have

$$(1.2) \quad \|D^2u\|_{L^p(\Omega')} \leq C(\|u\|_{L^p(\Omega)} + \|f\|_{L^p(\Omega)}),$$

where C is a constant depending only on $p, \lambda, \Lambda, n, \text{dist}(\Omega', \partial\Omega)$ and the modulus of continuity of $A(x)$. The continuity assumption on the coefficient matrix is essential when $n \geq 3$. Indeed, it is shown in [U] and [PT] that if $A(x)$ satisfies (1.1) and is merely measurable, then (1.2) is false for $p \geq 1$. However, it is proved in [E, L] that estimates for second derivatives that do not depend on the continuity of $A(x)$ do hold when $p > 0$ is sufficiently small.

L^p -estimates for second derivatives of solutions to fully nonlinear uniformly elliptic equations of the form $F(D^2u, x) = f(x)$ were studied by Caffarelli [C1].

July 1, 2018.

CG gratefully acknowledges the support provided by NSF grant DMS-0901430.

TN gratefully acknowledges the support provided by NSF grant DMS-0901449.

In this fundamental work, he established Calderón-Zygmund type interior $W^{2,p}$ estimates for viscosity solutions under the assumptions that $F(D^2u, x)$ is suitably close to $F(D^2u, 0)$, and solutions to the frozen equation $F(D^2u, 0) = 0$ admit interior $C^{1,1}$ estimates. For more details and related results to those of Caffarelli, we refer to [CC, Chapter 7] and [Es, E, Sw, WL]. By extending further his perturbation method in [C1], Caffarelli [C3] was able to derive interior $W^{2,p}$ estimates for convex solutions to the Monge–Ampère equation $\det D^2\phi = g(x)$ under the optimal condition that g is continuous and bounded away from zero and infinity (see also [G, Chapter 6], [H, dPF] and the recent corresponding boundary estimates in [S2]).

In this paper we consider the linearized Monge–Ampère equation. Let $\Omega \subset \mathbb{R}^n$ be a normalized convex domain and $\phi \in C(\overline{\Omega})$ be a convex function satisfying $\lambda \leq \det D^2\phi = g(x) \leq \Lambda$ in Ω and $\phi = 0$ on $\partial\Omega$. The linearized Monge–Ampère equation corresponding to ϕ is

$$(1.3) \quad \mathcal{L}_\phi u := \text{trace}(\Phi D^2 u) = f(x) \quad \text{in } \Omega$$

where $\Phi := (\det D^2\phi) (D^2\phi)^{-1}$ is the matrix of cofactors of $D^2\phi$. We note that \mathcal{L}_ϕ is both a non divergence and divergence differential operator which is degenerate elliptic, that is, the matrix $\Phi(x)$ is positive semi-definite and does not satisfy (1.1). The equation (1.3) is of great importance as it appears in a number of problems. For example, it appears in affine differential geometry in the solution of the affine Bernstein problem ([T, TW1, TW2, TW3, TW4]), and in the Aubreu’s equation arising in the differential geometry of toric varieties ([D1, D2, D3, D4, Z1, Z2]). In addition, the equation appears in fluid mechanics in the semigeostrophic system which is an approximation to the incompressible Euler equation and is used in meteorology to study atmospheric flows ([CNP, Lo]). The linearized Monge–Ampère equation was first studied by Caffarelli and Gutiérrez in [CG2] where it is proved that nonnegative solutions to $\mathcal{L}_\phi u = 0$ satisfy a uniform Harnack’s inequality yielding, in particular, interior Hölder continuity of solutions. By using these interior Hölder estimates and perturbation arguments, we recently established in [GN] Cordes–Nirenberg type interior $C^{1,\alpha}$ estimates for solutions to (1.3).

The purpose in this paper is to study the L^p integrability of second derivatives of solutions to the equation (1.3). A previous result in this direction is proved by Gutiérrez and Tournier in [GT]: for any domain $\Omega' \Subset \Omega$, there exist $p > 0$ small and $C > 0$ depending only on λ, Λ, n and $\text{dist}(\Omega', \Omega)$ such that

$$(1.4) \quad \|D^2 u\|_{L^p(\Omega')} \leq C(\|u\|_{L^\infty(\Omega)} + \|f\|_{L^n(\Omega)})$$

for all solutions $u \in C^2(\Omega)$ of (1.3). Notice that since $\mathcal{L}_\phi \phi = n \det D^2\phi = ng(x)$, it follows from Wang’s counterexample [W] that (1.4) is false for any $p > 1$. In fact, if we hope the estimate (1.4) to hold for large values of p , one needs to assume in addition that $g \in C(\Omega)$, see [GT, Section 8] for more details. In light of this, it was conjectured in [GT] that the L^p -integrability of the second derivatives of u in (1.3)

improves when $\det D^2\phi$ gets closer to one; in other words, if $1-\epsilon \leq \det D^2\phi \leq 1+\epsilon$, then the exponent $p = p(\epsilon)$ in (1.4) satisfies $p(\epsilon) \rightarrow +\infty$ as $\epsilon \rightarrow 0$.

In this article we solve the above conjecture in the affirmative, Theorem 4.6. As a consequence, we obtain the following main result of the paper.

Theorem 1.1. *Let Ω be a normalized convex domain and $g \in C(\Omega)$ with $0 < \lambda \leq g(x) \leq \Lambda$. Suppose $u \in W_{loc}^{2,n}(\Omega)$ is a solution of $\mathcal{L}_\phi u = f$ in Ω , where $\phi \in C(\overline{\Omega})$ is a convex function satisfying $\det D^2\phi = g$ in Ω and $\phi = 0$ on $\partial\Omega$. Let $\Omega' \Subset \Omega$, $p > 1$ and $\max\{n, p\} < q < \infty$. Then there exists $C > 0$ depending only on $p, q, \lambda, \Lambda, n, \text{dist}(\Omega', \partial\Omega)$ and the modulus of continuity of g such that*

$$(1.5) \quad \|D^2u\|_{L^p(\Omega')} \leq C \left(\|u\|_{L^\infty(\Omega)} + \|f\|_{L^q(\Omega)} \right).$$

The conditions on the Monge-Ampère measure $\det D^2\phi$ are sharp and the constant in (1.5) depends on $\det D^2\phi$ and not on the maximum or minimum of eigenvalues of $D^2\phi$. Our result can be viewed as a degenerate counterpart of the classical Calderón-Zygmund estimates (1.2) for linear uniformly elliptic equations in non divergence form, and Caffarelli's interior $W^{2,p}$ estimates [C1, Theorem 1], [CC, Chapter 7] for fully nonlinear uniformly elliptic equations.

In order to address the lack of uniform ellipticity of the linearized Monge-Ampère operator, we follow the strategy in [CG2] by working with sections of solutions to the Monge-Ampère equation. The role that the sections play in our analysis is similar to that of Euclidean balls in the theory of uniformly elliptic equations. In addition, to measure the degree of regularity of the solution we introduce the sets $G_M(u, \Omega)$ where the solution u is touched by tangent paraboloids, see Definition 2.6. In contrast with [GT, Definition 3.5], the sets $G_M(u, \Omega)$ are now invariant by affine transformations. We note that unlike the theory in [CC, Chapter 7], where the standard Euclidean distance is used, our tangent paraboloids are defined with respect to a quasi distance induced by the solution ϕ of the Monge-Ampère equation. With this new definition, our first step is to derive rough density estimates for the sets $G_M(u, \Omega)$ which are achieved by following the method in [GT]. The next crucial step in solving the conjecture is to accelerate the initial density estimates. To make this breakthrough, we use a key idea introduced in [GN], that is, to compare solutions of two different linearized Monge-Ampère equations. Precisely, we compare solutions of $\mathcal{L}_\phi u = f$ with solutions of $\mathcal{L}_w h = 0$, having the same Dirichlet boundary data, where w is the solution of the Monge-Ampère equation $\det D^2w = 1$ in Ω and $w = 0$ on $\partial\Omega$. It is also important to know that the coefficient matrices of two different linearized equations are close in L^p -norm when the determinants of the corresponding convex functions are close in L^∞ -norm. This is given in our recent work [GN]. These two comparison results allow us to estimate explicitly $\|u - h\|_{L^\infty}$ in terms of $\|\det D^2\phi - 1\|_{L^\infty}$, and by using this approximation we can perform the acceleration process to obtain the necessary density estimates for the sets $G_M(u, \Omega)$. Finally and to conclude the proof of the conjecture, all these estimates permit us to use

the covering theorems for sections of solutions to the Monge-Ampère equation proved in [CG1, CG2].

To give more perspective, we mention the following recent work for the linearized Monge-Ampère equation: Sobolev type inequalities associated to the linearized operator \mathcal{L}_ϕ (Tian and Wang [TiW]), Liouville property for solutions to $\mathcal{L}_\phi u = 0$ in \mathbb{R}^2 (Savin [S1]) and boundary $C^{1,\alpha}$ estimates for $\mathcal{L}_\phi u = f$ and its applications (Le and Savin [LS1, LS2]).

The paper is organized as follows. Section 2 is devoted to preliminary results for solutions ϕ to the Monge-Ampère equation that will be used later. We also introduce there a quasi metric and the sets $G_M(u, \Omega)$ where the solution u to the linearized equation is touched by tangent paraboloids associated to the quasi distance. In Section 3 we establish density estimates for the set $G_M(u, \Omega)$ and use them to derive the initial power decay for the distribution function giving small integrability of D^2u . Finally, Section 4 contains the main estimates in the paper showing how the integrability improves when $\det D^2\phi$ gets closer to one.

2. PRELIMINARY RESULTS

2.1. Some properties for the Monge-Ampère equation. Given an open convex set $\Omega \subset \mathbb{R}^n$ and a function $\phi \in C(\Omega)$, $\partial\phi$ denotes the sub differential of ϕ . The Monge-Ampère measure associated with ϕ is defined by $M\phi(E) := |\partial\phi(E)|$, for all Borel subsets $E \subset \Omega$. The convex set Ω is called a *normalized convex domain* if $B_1(0) \subset \Omega \subset B_n(0)$. Here $B_R(y)$ denotes the Euclidean ball with radius R centered at y . Observe that by Fritz John's lemma, every bounded convex domain with non empty interior can be normalized, i.e., there is an invertible affine transformation T with $B_1(0) \subset T(S) \subset B_n(0)$. A *section of a convex function* $\phi \in C^1(\Omega)$ centered at \bar{x} and with height t is defined by

$$S_\phi(\bar{x}, t) = \{x \in \Omega : \phi(x) < \phi(\bar{x}) + \nabla\phi(\bar{x}) \cdot (x - \bar{x}) + t\}.$$

If $\phi = 0$ on $\partial\Omega$, then for $0 < \alpha < 1$ we set

$$(2.6) \quad \Omega_\alpha = \{x \in \Omega : \phi(x) < (1 - \alpha) \min_\Omega \phi\},$$

and notice that Ω_α is a section of ϕ at the minimum of ϕ , i.e., $\Omega_\alpha = S_\phi(x_0, -\alpha\phi(x_0))$ where $x_0 \in \Omega$ is such that $\min_\Omega \phi = \phi(x_0)$. We are going to list some basic properties related to sections that will be used later. All results in this subsection hold under the assumption:

(H) Ω is a normalized convex domain and $\phi \in C(\overline{\Omega})$ is a convex function such that

$$\lambda \leq M\phi \leq \Lambda \quad \text{in } \Omega \quad \text{and} \quad \phi = 0 \quad \text{on } \partial\Omega.$$

It is known from the works of Caffarelli [C2, C4] that ϕ is strictly convex and $C^{1,\alpha}$ in the interior of Ω . Moreover, we have the following lemma from [GH] (see [G, Theorem 3.3.10]).

Lemma 2.1. *Let $0 < \alpha < \beta < 1$. Then for any $x \in \Omega_\alpha$, we have $S_\phi(x, C_0(\beta - \alpha)^\gamma) \subset \Omega_\beta$ for some C_0 and γ depending only on n, λ and Λ . Consequently, there exists $\eta = \eta(\alpha, n, \lambda, \Lambda) > 0$ such that $S_\phi(x, t) \Subset \Omega$ for all $x \in \Omega_\alpha$ and $t \leq \eta$.*

We now state a result about strong type $p - p$ estimates for the maximal function with respect to sections. For that, let us fix $0 < \alpha_0 < 1$ and take $\eta_0 = \eta_0(\alpha_0, n, \lambda, \Lambda)$ be the corresponding positive constant given by Lemma 2.1.

Theorem 2.2. *Let $\mu := M\phi$ and define*

$$\mathcal{M}_\mu(f)(x) := \sup_{t \leq \eta_0/2} \frac{1}{\mu(S_\phi(x, t))} \int_{S_\phi(x, t)} |f(y)| d\mu(y) \quad \forall x \in \Omega_{\alpha_0}.$$

Then for any $1 < p < \infty$, there exists a constant C depending on p, n, λ and Λ such that

$$\left(\int_{\Omega_{\alpha_0}} |\mathcal{M}_\mu(f)(x)|^p d\mu(x) \right)^{\frac{1}{p}} \leq C \left(\int_{\Omega} |f(y)|^p d\mu(y) \right)^{\frac{1}{p}}.$$

Notice that it is known from [CG1] and [GT, Theorem 2.9] that \mathcal{M}_μ is of weak type $1 - 1$. This together with the obvious inequality $\|\mathcal{M}_\mu(f)\|_{L^\infty(\Omega_{\alpha_0})} \leq \|f\|_{L^\infty(\Omega)}$ and the Marcinkiewicz interpolation lemma (see Theorem 5 in [St, Page 21]) yields the strong type $p - p$ estimate in Theorem 2.2. The next lemma is a slight modification of [G, Lemma 6.2.1].

Lemma 2.3. *There exist $c = c(n, \lambda, \Lambda) > 0$ and $\delta_0 = \delta_0(\alpha_0, n, \lambda, \Lambda) > 0$ such that if $x_0 \in \Omega_{\alpha_0}$ and $\phi(x) \geq \phi(x_0) + \nabla\phi(x_0) \cdot (x - x_0) + \sigma|x - x_0|^2 \quad \forall x \in \Omega$, then*

$$\phi(x) \leq \phi(x_0) + \nabla\phi(x_0) \cdot (x - x_0) + \frac{1}{c^2\sigma^{n-1}}|x - x_0|^2 \quad \text{for all } |x - x_0| \leq \delta_0.$$

Proof. Let $u(x) := \phi(x) - \phi(x_0) - \nabla\phi(x_0) \cdot (x - x_0)$. Then by the proof of Lemma 6.2.1 in [G], we have $u(x) \leq C(n, \lambda, \Lambda)\sigma^{-n+1}|x - x_0|^2$ for all $x \in \Omega$ satisfying $u(x) \leq \eta_0$. Next it follows from Aleksandrov's maximum principle and [G, Proposition 3.2.3] that $\text{dist}(\Omega_{\alpha_0}, \partial\Omega) \geq c(n, \lambda, \Lambda)(1 - \alpha_0)^n =: d_{\alpha_0}$. Moreover if $\text{dist}(x, \partial\Omega) \geq d_{\alpha_0}/2$, then by using [G, Lemma 3.2.1] we obtain

$$u(x) \leq |\nabla\phi(\xi) - \nabla\phi(x_0)| |x - x_0| \leq \frac{C(n, \lambda, \Lambda)}{d_{\alpha_0}} |x - x_0|$$

where ξ is some point on the segment joining x_0 and x . Therefore there exists $\delta_0 > 0$ depending only on α_0, n, λ and Λ such that $u(x) \leq \eta_0$ whenever $|x - x_0| \leq \delta_0$. \square

The above lemma together with Lemma 6.2.2 in [G] gives:

Lemma 2.4. *Given $0 < \alpha \leq \alpha_0$ and $\gamma > 0$, we define*

$$(2.7) \quad D_\gamma^\alpha = \left\{ x \in \Omega_\alpha : S_\phi(x, t) \subset B_{\gamma\sqrt{t}}(x), \quad \forall t \leq \eta_0 \right\}.$$

Then there exist $c = c(n, \lambda, \Lambda) > 0$ and $\delta_0 = \delta_0(\alpha_0, n, \lambda, \Lambda) > 0$ such that for any $\gamma > 0$ satisfying $(c\gamma)^{\frac{1}{n-1}} \geq \frac{\text{diam}(\Omega)}{\sqrt{\eta_0}}$, we have: if $\bar{x} \in D_{(c\gamma)^{\frac{1}{n-1}}}^\alpha$ then

$$\phi(x) - \phi(\bar{x}) - \nabla\phi(\bar{x}) \cdot (x - \bar{x}) \leq \gamma^2 |x - \bar{x}|^2 \quad \text{for all } |x - \bar{x}| \leq \delta_0.$$

Proof. Let c and δ_0 be given by Lemma 2.3, and take $\bar{x} \in D_{(c\gamma)^{\frac{1}{n-1}}}^\alpha$. Since $(c\gamma)^{\frac{1}{n-1}} \geq \frac{\text{diam}(\Omega)}{\sqrt{\eta_0}}$, we then have by [G, Lemma 6.2.2] that $\bar{x} \in \Omega_\alpha \cap A_{(c\gamma)^{\frac{1}{n-1}}}$, where

$$A_\sigma := \left\{ x_0 \in \Omega : \phi(x) \geq \phi(x_0) + \nabla\phi(x_0) \cdot (x - x_0) + \sigma |x - x_0|^2, \quad \forall x \in \Omega \right\}.$$

Therefore the conclusion of the lemma follows from Lemma 2.3. \square

2.2. Tangent paraboloids and power decay for the Monge-Ampère equation.

In this subsection we recall the quasi distance given by the convex function ϕ and then use it to define the sets where the solution u is touched from above and below by certain functions involving this quasi distance.

Definition 2.5. Let Ω be a bounded convex set in \mathbb{R}^n and $\phi \in C^1(\Omega)$ be a convex function. For any $x \in \Omega$ and $x_0 \in \Omega$, we define $d(x, x_0)$ by

$$d(x, x_0)^2 := \phi(x) - \phi(x_0) - \nabla\phi(x_0) \cdot (x - x_0).$$

Clearly $x \mapsto d(x, x_0)^2$ is a convex function on Ω . Since $d(x, x_0)^2$ is in general not equivalent to $|x - x_0|^2$, the following definition of "tangent paraboloids" has a nature different from the standard definition of tangent paraboloids for uniformly elliptic equations (see [CC]). It is however more suitable to exploit the degenerate structure of the solution ϕ to the Monge-Ampère equation.

Definition 2.6. Let Ω and ϕ be as in Definition 2.5. Then for $u \in C(\Omega)$ and $M > 0$, we define the sets

$$G_M^+(u, \Omega) = \left\{ \bar{x} \in \Omega : u \text{ is differentiable at } \bar{x} \text{ and } u(x) \leq u(\bar{x}) + \nabla u(\bar{x}) \cdot (x - \bar{x}) + M d(x, \bar{x})^2 \quad \forall x \in \Omega \right\};$$

$$G_M^-(u, \Omega) = \left\{ \bar{x} \in \Omega : u \text{ is differentiable at } \bar{x} \text{ and } u(x) \geq u(\bar{x}) + \nabla u(\bar{x}) \cdot (x - \bar{x}) - M d(x, \bar{x})^2 \quad \forall x \in \Omega \right\};$$

and $G_M(u, \Omega) := G_M^+(u, \Omega) \cap G_M^-(u, \Omega)$.

We note that local versions of Definition 2.5 and Definition 2.6 were introduced in [GT]. However, these definitions are not good enough for the purposes of this paper. The next observation is our starting point for deriving L^p -estimates for second derivatives of solutions to the linearized Monge-Ampère equation.

Lemma 2.7. Assume condition (H) holds. Let $0 < \alpha \leq \alpha_0$, $u \in C(\Omega)$ and

$$(2.8) \quad \Theta(u)(x) := \frac{1}{2} \Theta(u, B_{\delta_0}(x))(x) \quad \text{for } x \in \Omega_{\alpha_0}$$

where δ_0 is given by Lemma 2.4 and $\Theta(u, B_{\delta_0}(x))(x)$ is defined exactly as in [CC, Section 1.2]. Then for $\kappa > 1$, we have

$$(2.9) \quad \{x \in \Omega_\alpha : \Theta(u)(x) > \beta^\kappa\} \subset \left(\Omega_\alpha \setminus D_{(c\beta^{\frac{\kappa-1}{2}})^{\frac{1}{n-1}}}^\alpha \right) \cup \left(\Omega_\alpha \setminus G_\beta(u, \Omega) \right)$$

for any $\beta > 0$ satisfying $(c\beta^{\frac{\kappa-1}{2}})^{\frac{1}{n-1}} \geq \frac{\text{diam}(\Omega)}{\sqrt{\eta_0}}$ with $c = c(n, \lambda, \Lambda)$ is as in Lemma 2.4.

Proof. Let $\gamma := \beta^{\frac{\kappa-1}{2}}$. If $\bar{x} \in D_{(c\gamma)^{\frac{1}{n-1}}}^\alpha \cap G_\beta(u, \Omega)$, then

$$-\beta d(x, \bar{x})^2 \leq u(x) - u(\bar{x}) - \nabla u(\bar{x}) \cdot (x - \bar{x}) \leq \beta d(x, \bar{x})^2$$

for each $x \in \Omega$. Since $\bar{x} \in D_{(c\gamma)^{\frac{1}{n-1}}}^\alpha$, this together with Lemma 2.4 yields

$$-\beta\gamma^2|x - \bar{x}|^2 \leq u(x) - u(\bar{x}) - \nabla u(\bar{x}) \cdot (x - \bar{x}) \leq \beta\gamma^2|x - \bar{x}|^2$$

for all $|x - \bar{x}| \leq \delta_0$, and so $\Theta(u, B_{\delta_0}(x))(\bar{x}) \leq 2\beta\gamma^2 = 2\beta^\kappa$. Thus we have proved that

$$D_{(c\gamma)^{\frac{1}{n-1}}}^\alpha \cap G_\beta(u, \Omega) \subset \{x \in \Omega_\alpha : \Theta(u)(x) \leq \beta^\kappa\}$$

and the lemma follows by taking complements. \square

In order to derive interior $W^{2,p}$ estimates for solutions u to the linearized Monge-Ampère equation, we will need to estimate the distribution function $F(\beta) := |\{x \in \Omega_\alpha : \Theta(u)(x) > \beta^\kappa\}|$ for some suitable choice of $\kappa > 1$. It follows from Lemma 2.7 that this can be done if one can get appropriate fall off estimates for $F_1(\beta) := |\Omega_\alpha \setminus D_{(c\beta^{\frac{\kappa-1}{2}})^{\frac{1}{n-1}}}^\alpha|$ and $F_2(\beta) := |\Omega_\alpha \setminus G_\beta(u, \Omega)|$ when β is large. Notice that since the function $F_1(\beta)$ involves only the solution ϕ of the Monge-Ampère equation, its decay estimate has been established by Caffarelli in the fundamental work [C3]. We reformulate his estimate in the following theorem.

Theorem 2.8. *Let Ω be a normalized convex domain and $\phi \in C(\overline{\Omega})$ be a convex function satisfying $1 - \epsilon \leq \det D^2\phi \leq 1 + \epsilon$ in Ω and $\phi = 0$ on $\partial\Omega$, where $0 < \epsilon < 1/2$. Then for any $0 < \alpha < 1$, there exists a positive constant M depending only on α and n such that*

$$(2.10) \quad |\Omega_\alpha \setminus D_s^\alpha| \leq \frac{|\Omega|}{(C_n\epsilon)^2} s \frac{\ln \sqrt{C_n\epsilon}}{\ln M} \quad \text{for all } s \geq M.$$

Proof. This theorem is obtained by iterating [G, Theorem 6.3.2]. Indeed, let $\alpha_0 := \frac{\alpha+1}{2}$ and let $M = M(\alpha_0, n)$ and $p_0 = p_0(\alpha_0, n)$ be the positive constants given by that theorem. By taking if necessary an even bigger constant, we can assume that M is large so that $\alpha_0 - \sum_{j=1}^{\infty} M^{-(j+1)p_0} \geq 2\alpha_0 - 1$ and the statement of [G, Theorem 6.3.2] holds for all $\lambda \geq M$. We then begin the iteration with $\lambda = M$ and let $\alpha_1 = \alpha_0 - M^{-2p_0}$. We get from [G, Theorem 6.3.2] that

$$|\Omega_{\alpha_1} \setminus D_{M^2}^{\alpha_1}| \leq \sqrt{C_n\epsilon} |\Omega_{\alpha_0} \setminus D_M^{\alpha_0}|.$$

If $\lambda = M^2$ and $\alpha_2 = \alpha_1 - M^{-3p_0}$, then

$$|\Omega_{\alpha_2} \setminus D_{M^3}^{\alpha_2}| \leq \sqrt{C_n \epsilon} |\Omega_{\alpha_1} \setminus D_{M^2}^{\alpha_1}| \leq \left(\sqrt{C_n \epsilon}\right)^2 |\Omega_{\alpha_0} \setminus D_M^{\alpha_0}|.$$

Continuing in this way we let $\alpha_{k-1} = \alpha_{k-2} - M^{-kp_0} = \alpha_0 - \sum_{j=1}^{k-1} M^{-(j+1)p_0}$ and obtain

$$|\Omega_{\alpha_{k-1}} \setminus D_{M^k}^{\alpha_{k-1}}| \leq \left(\sqrt{C_n \epsilon}\right)^{k-1} |\Omega_{\alpha_0} \setminus D_M^{\alpha_0}|.$$

Since by our choice of M , $\alpha_{k-1} \geq \alpha_0 - \sum_{j=1}^{\infty} M^{-(j+1)p_0} \geq 2\alpha_0 - 1$, it is easy to see that $\Omega_{2\alpha_0-1} \setminus D_{M^k}^{2\alpha_0-1} \subset \Omega_{\alpha_{k-1}} \setminus D_{M^k}^{\alpha_{k-1}}$. Therefore, we have

$$|\Omega_{\alpha} \setminus D_{M^k}^{\alpha}| = |\Omega_{2\alpha_0-1} \setminus D_{M^k}^{2\alpha_0-1}| \leq \left(\sqrt{C_n \epsilon}\right)^{k-1} |\Omega_{\alpha_0} \setminus D_M^{\alpha_0}| \quad \text{for } k = 1, 2, \dots$$

Now for each $s \geq M$, let us pick k such that $M^k \leq s \leq M^{k+1}$. Then $D_{M^k}^{\alpha} \subset D_s^{\alpha} \subset D_{M^{k+1}}^{\alpha}$ and $k \leq \log_M s \leq k+1$. So

$$|\Omega_{\alpha} \setminus D_s^{\alpha}| \leq \frac{|\Omega_{\alpha_0}|}{(C_n \epsilon)^2} \left(\sqrt{C_n \epsilon}\right)^{k+1} \leq \frac{|\Omega_{\alpha_0}|}{(C_n \epsilon)^2} \left(\sqrt{C_n \epsilon}\right)^{\log_M s} = \frac{|\Omega_{\alpha_0}|}{(C_n \epsilon)^2} s^{\frac{\ln \sqrt{C_n \epsilon}}{\ln M}}.$$

□

3. L^δ ESTIMATES FOR SECOND DERIVATIVES

In this section we prove two density lemmas and then use them to prove a small power decay of $\mu(\Omega_{\alpha} \setminus G_{\beta}(u, \Omega))$ for β large. Observe that the density estimates established in [GT] are not good enough for our purpose since a different definition of the sets $G_{\beta}(u)$ was introduced there. In [GT, Definition 3.5] the "tangent paraboloid" is assumed to lie below or above u in a specific neighborhood depending on β of the touching point. Such definition is not invariant under normalization and so not suitable for the acceleration process we consider later in Section 4. In this paper, we employ a global definition, Definition 2.6, and we are still able to obtain similar estimates as in [GT] by modifying their arguments. For clarity, in the next subsection we give complete proofs of these estimates that are technically simpler than the ones in [GT]. The following lemma is an extension of [GT, Lemma 3.1] which allows us to work with strong solutions in $W_{loc}^{2,n}(\Omega)$ instead of classical solutions.

Lemma 3.1. *Let $\Omega \subset \mathbb{R}^n$ be open and $u, \phi \in W_{loc}^{2,n}(\Omega)$ be such that $\det D^2 \phi(x) > 0$ for almost every x in Ω . Let $w = u + \phi$. Then for any Borel set $E \subset \Omega$, we have*

$$(3.11) \quad Mw(E) \leq \frac{1}{n^n} \int_{E \cap C} \left(\left(\frac{\text{trace}(\Phi(x) D^2 u(x))}{\det D^2 \phi(x)} + n \right)^+ \right)^n \det D^2 \phi(x) dx$$

where $\Phi(x)$ is the matrix of cofactors of $D^2 \phi(x)$ and $C := \{x \in \Omega : w(x) = \Gamma(w)(x)\}$ with $\Gamma(w)$ is the convex envelope of w in Ω .

Proof. Notice that the Sobolev embedding theorem guarantees that functions in $W_{loc}^{2,n}(\Omega)$ are continuous in Ω . We first claim that

$$(3.12) \quad Mw(F) \leq \int_F |\det D^2 w(x)| dx \quad \text{for all Borel sets } F \subset \Omega.$$

It is well known that (3.12) holds if $w \in C^2(\Omega)$. For general $w \in W_{loc}^{2,n}(\Omega)$, let $\{w_m\}$ be a sequence of functions in $C^2(\Omega)$ converging to w in the sense of $W_{loc}^{2,n}(\Omega)$. Let $U \subset \Omega$ be open and $K \subset U$ be compact. Then $K \subset U \cap \Omega_\epsilon$ for all $\epsilon > 0$ sufficiently small, where $\Omega_\epsilon := \{x \in \Omega : \text{dist}(x, \partial\Omega) > \epsilon\}$. Since $Mw_m(U \cap \Omega_\epsilon) \leq \int_{U \cap \Omega_\epsilon} |\det D^2 w_m(x)| dx$, we get

$$\limsup_{m \rightarrow \infty} Mw_m(U \cap \Omega_\epsilon) \leq \limsup_{m \rightarrow \infty} \int_{U \cap \Omega_\epsilon} |\det D^2 w_m - \det D^2 w| dx + \int_{U \cap \Omega_\epsilon} |\det D^2 w| dx.$$

Since the first term on the right hand side is clearly zero and the measures Mw_m converge to the measure Mw weakly, it follows by taking $\epsilon > 0$ small enough that $Mw(K) \leq \int_U |\det D^2 w| dx$. Consequently,

$$(3.13) \quad Mw(U) \leq \int_U |\det D^2 w(x)| dx$$

by the regularity of the measure Mw . Because (3.13) is true for any open set $U \subset \Omega$, we once again use the regularity of the measures to infer that the claim (3.12) holds.

Now let $E \subset \Omega$ be an arbitrary Borel set. It is clear that $\partial w(E) = \partial w(E \cap C)$ and so by using (3.12) and the fact $D^2 w(x) \geq 0$ for almost every x in C we obtain

$$Mw(E) = Mw(E \cap C) \leq \int_{E \cap C} \det D^2 w(x) dx$$

and the estimate (3.11) follows by a calculation from [GT, Lemma 3.1]. \square

Throughout this paper we always work with strong solutions, in the Sobolev space $W_{loc}^{2,n}(\Omega)$, of the linearized Monge-Ampère equation. That is, the equation $\mathcal{L}_\phi u = f$ in Ω is interpreted in the almost everywhere sense in Ω .

3.1. Initial density estimates.

Lemma 3.2. *Let U be a normalized convex domain and Ω be a bounded convex set such that $U \subset \Omega$. Let $\phi \in C^1(\Omega) \cap W_{loc}^{2,n}(U)$ be a convex function satisfying $\lambda \leq \det D^2 \phi \leq \Lambda$ in U . Suppose $u \in C(\Omega) \cap W_{loc}^{2,n}(U) \cap C^1(U)$, $0 \leq u \leq 1$ in Ω and $\mathcal{L}_\phi u = f$ in U . Then for each $\epsilon > 0$ there exists $\eta(\epsilon, n, \lambda, \Lambda) > 0$ such that for any $\eta \leq \eta(\epsilon, n, \lambda, \Lambda)$, we have*

$$\mu\left(G_{\frac{1}{\eta_0}}^-(u, \Omega) \cap S_\phi(x_0, t_0)\right) \geq \left[(1 - \epsilon)^{\frac{1}{n}} - C\eta t_0 \left(\int_{S_\phi(x_0, t_0)} \left| \frac{f}{\det D^2 \phi} \right|^n d\mu \right)^{\frac{1}{n}} \right]^n \mu(S_\phi(x_0, t_0))$$

for all sections $S_\phi(x_0, t_0) \Subset U$. Here $\mu := M\phi$ and C depends only on n, λ and Λ .

Proof. Let T normalize the section $S_\phi(x_0, t_0)$. For $y \in T(\Omega)$, set

$$\tilde{\phi}(y) = \frac{1}{t_0} \left[\phi(T^{-1}y) - \phi(x_0) - \nabla\phi(x_0) \cdot (T^{-1}y - x_0) - t_0 \right] \quad \text{and} \quad \tilde{u}(y) = u(T^{-1}y).$$

We have that $\tilde{\Omega} := T(S_\phi(x_0, t_0))$ is normalized and $\tilde{\phi} = 0$ on $\partial\tilde{\Omega}$. Also, it follows from [GN, Lemma 2.3] that $\lambda' \leq \det D^2\tilde{\phi} \leq \Lambda'$ in $\tilde{\Omega}$, where λ' and Λ' depend only on n , λ and Λ . By Lemma 2.1, for each $0 < \alpha < 1$ there exists $\eta(\alpha) = \eta(\alpha, n, \lambda, \Lambda)$ such that if $\bar{y} \in \tilde{\Omega}_\alpha$, then $S_{\tilde{\phi}}(\bar{y}, \eta(\alpha)) \Subset \tilde{\Omega}$. Therefore if $\bar{y} \in \tilde{\Omega}_\alpha$, then

$$(3.14) \quad \tilde{\phi}(\bar{y}) + \nabla\tilde{\phi}(\bar{y}) \cdot (y - \bar{y}) + \eta(\alpha) < 0 \quad \text{for all } y \in \overline{\tilde{\Omega}}.$$

Define $w_\alpha(y) = \eta(\alpha)\tilde{u}(y) + \tilde{\phi}(y)$. Let γ_α be the convex envelope of w_α in $\tilde{\Omega}$ and

$$C_\alpha = \{y \in \tilde{\Omega} : w_\alpha(y) = \gamma_\alpha(y),$$

and $\exists \ell_y$ supporting hyperplane to γ_α at y with $\ell_y < 0$ in $\tilde{\Omega}\}$.

Claim 1. $\nabla\tilde{\phi}(\tilde{\Omega}_\alpha) = \nabla(\tilde{\phi} + \eta(\alpha))(\tilde{\Omega}_\alpha) \subset \nabla w_\alpha(C_\alpha)$.

To prove this claim, note first that $\tilde{\phi} + \eta(\alpha) \geq w_\alpha$ in $\tilde{\Omega}$ and $w_\alpha \geq 0$ on $\partial\tilde{\Omega}$. If $\bar{y} \in \tilde{\Omega}_\alpha$, then by (3.14) we know that the supporting plane $z = \tilde{\phi}(\bar{y}) + \nabla\tilde{\phi}(\bar{y}) \cdot (y - \bar{y}) + \eta(\alpha)$ to $\tilde{\phi} + \eta(\alpha)$ at \bar{y} has the property: $z < 0$ on $\partial\tilde{\Omega}$. Therefore, if we slide it down, then it must become a supporting plane to w_α at some point $y^* \in \tilde{\Omega}$ (say ℓ_{y^*}). Since $z < 0$ in $\tilde{\Omega}$, so is ℓ_{y^*} and hence $y^* \in C_\alpha$. Thus $\nabla\tilde{\phi}(\bar{y}) \in \nabla w_\alpha(C_\alpha)$ as desired.

Claim 2. $C_\alpha \subset T(G_{1/(t_0\eta(\alpha))}^-(u, \Omega) \cap S_\phi(x_0, t_0))$, for every $0 < \alpha < 1$.

Proof of Claim 2. Let $\bar{y} \in C_\alpha$. Then $\bar{y} = T\bar{x}$ for some $\bar{x} \in S_\phi(x_0, t_0)$ and

$$(3.15) \quad \eta(\alpha)\tilde{u}(y) + \tilde{\phi}(y) \geq \ell(y) \quad \forall y \in T(S_\phi(x_0, t_0))$$

with equality at $y = \bar{y}$, for some ℓ affine with $\ell < 0$ in $T(S_\phi(x_0, t_0))$. As $\phi(x) = \phi(\bar{x}) + \nabla\phi(\bar{x}) \cdot (x - \bar{x}) + d(x, \bar{x})^2$ in Ω , we have

$$\tilde{\phi}(y) = \tilde{\phi}(\bar{y}) + \nabla\tilde{\phi}(\bar{y}) \cdot (y - \bar{y}) + t_0^{-1} d(T^{-1}y, T^{-1}\bar{y})^2 =: \tilde{\ell}_{\bar{y}}(y) + t_0^{-1} d(T^{-1}y, T^{-1}\bar{y})^2$$

for all $y \in T(\Omega)$. This together with (3.15) gives

$$\eta(\alpha)\tilde{u}(y) \geq \ell(y) - \tilde{\ell}_{\bar{y}}(y) - t_0^{-1} d(T^{-1}y, T^{-1}\bar{y})^2 =: g(y) \quad \forall y \in T(S_\phi(x_0, t_0))$$

with equality at $y = \bar{y}$. Assume for a moment that

$$(3.16) \quad 0 \geq g(y) \quad \text{for all } y \in T(\Omega) \setminus T(S_\phi(x_0, t_0)).$$

Since $u \geq 0$ in Ω , we then obtain

$$(3.17) \quad \eta(\alpha)\tilde{u}(y) \geq g(y) \quad \forall y \in T(\Omega).$$

To see (3.16), let $B := \{y \in T(\Omega) : g(y) \geq 0\}$. Note that $\bar{y} \in B$. Also $B \cap \partial T(S_\phi(x_0, t_0)) = \emptyset$ because if $y \in \partial T(S_\phi(x_0, t_0))$ then $\ell(y) < 0$ and so $g(y) < -\tilde{\ell}_{\bar{y}}(y) - t_0^{-1} d(T^{-1}y, T^{-1}\bar{y})^2 = -\tilde{\phi}(y) = 0$. Moreover, B is connected as g is concave. Hence $B \subset T(S_\phi(x_0, t_0))$ implying (3.16).

Since ℓ is a supporting hyperplane to $\eta(\alpha)\tilde{u}(y) + \tilde{\phi}(y)$ at \bar{y} , and $u, \phi \in C^1(U)$, it follows from (3.17) that

$$\tilde{u}(y) \geq \tilde{u}(\bar{y}) + \nabla \tilde{u}(\bar{y}) \cdot (y - \bar{y}) - \frac{1}{t_0 \eta(\alpha)} d(T^{-1}y, T^{-1}\bar{y})^2, \quad \forall y \in T(\Omega).$$

Thus we have proved that

$$C_\alpha \subset \{\bar{y} \in T(S_\phi(x_0, t_0)) : \tilde{u}(y) \geq \tilde{u}(\bar{y}) + \nabla \tilde{u}(\bar{y}) \cdot (y - \bar{y}) - \frac{1}{t_0 \eta(\alpha)} d(T^{-1}y, T^{-1}\bar{y})^2 \forall y \in T(\Omega)\}$$

yielding Claim 2 because $\tilde{u}(y) = u(T^{-1}y)$.

Now let $\tilde{\Phi}(y) := (D^2\tilde{\phi}(y))^{-1} \det D^2\tilde{\phi}(y)$. Then as $D^2\tilde{\phi}(y) = t_0^{-1}(T^{-1})^t D^2\phi(T^{-1}y) T^{-1}$ and $D^2\tilde{u}(y) = (T^{-1})^t D^2u(T^{-1}y) T^{-1}$, we get

$$\text{trace}(\tilde{\Phi}(y) D^2\tilde{u}(y)) = \frac{t_0}{t_0^n |\det T|^2} \text{trace}(\Phi(T^{-1}y) D^2u(T^{-1}y)) = \frac{t_0 f(T^{-1}y)}{t_0^n |\det T|^2} \quad \text{in } \tilde{\Omega}.$$

Therefore by applying Lemma 3.1 with $\Omega \rightsquigarrow \tilde{\Omega}$, $u \rightsquigarrow \eta(\alpha)\tilde{u}$, $\phi \rightsquigarrow \tilde{\phi}$, $E = C_\alpha$ and using Claim 1 and the fact $t_0^n |\det T|^2 \approx 1$, we obtain

$$\int_{\tilde{\Omega}_\alpha} \det D^2\tilde{\phi}(y) dy \leq \frac{1}{n^n} \int_{C_\alpha} \left(\frac{C\eta(\alpha) t_0 |f(T^{-1}y)|}{\det D^2\tilde{\phi}(y)} + n \right)^n \det D^2\tilde{\phi}(y) dy.$$

Since $C_\alpha \subset T(G_{1/(t_0\eta(\alpha))}^-(u, \Omega) \cap S_\phi(x_0, t_0))$ by Claim 2, $\det D^2\tilde{\phi}(y) = \det D^2\phi(T^{-1}y)$ and $\tilde{\Omega}_\alpha = T(S_\phi(x_0, \alpha t_0))$, the above inequality implies

$$\int_{S_\phi(x_0, \alpha t_0)} \det D^2\phi(x) dx \leq \frac{1}{n^n} \int_{G_{1/(t_0\eta(\alpha))}^-(u, \Omega) \cap S_\phi(x_0, t_0)} \left(\frac{C\eta(\alpha) t_0 |f(x)|}{\det D^2\phi(x)} + n \right)^n \det D^2\phi(x) dx.$$

We then infer from Minkowski's inequality and $\mu = M\phi$ that

$$\begin{aligned} & \mu(S_\phi(x_0, \alpha t_0))^{\frac{1}{n}} \\ & \leq \frac{C\eta(\alpha) t_0}{n} \left(\int_{S_\phi(x_0, t_0)} \left| \frac{f}{\det D^2\phi} \right|^n d\mu \right)^{\frac{1}{n}} \mu(S_\phi(x_0, t_0))^{\frac{1}{n}} + \mu(G_{1/(t_0\eta(\alpha))}^-(u, \Omega) \cap S_\phi(x_0, t_0))^{\frac{1}{n}}. \end{aligned}$$

Note that this inequality holds for any $\eta \leq \eta(\alpha)$. Given $\epsilon > 0$ there exists $\alpha = \alpha(\epsilon)$ sufficiently close to one such that $(1 - \epsilon)\mu(S_\phi(x_0, t_0)) \leq \mu(S_\phi(x_0, \alpha t_0))$, which combined with the previous inequality yields the lemma for any $\eta \leq \eta(\alpha(\epsilon))$. \square

In the next lemma, we no longer require $0 \leq u \leq 1$ in Ω as in Lemma 3.2.

Lemma 3.3. *Let U be a normalized convex domain and Ω be a bounded convex set such that $U \subset \Omega$. Let $\phi \in C^1(\Omega) \cap W_{loc}^{2,n}(U)$ be a convex function satisfying $\lambda \leq \det D^2\phi \leq \Lambda$ in U . Suppose $u \in C(\Omega) \cap W_{loc}^{2,n}(U) \cap C^1(U)$ is a solution of $\mathcal{L}_\phi u = f$ in U . Then for each*

$\epsilon > 0$ there exists $\eta(\epsilon, n, \lambda, \Lambda) > 0$ such that if $S_\phi(x_0, t_0) \Subset U$ and $S_\phi(x_0, t_0) \cap G_\gamma(u, \Omega)$ contains a point \bar{x} with $S_\phi(\bar{x}, \theta t_0) \Subset U$, then we have

$$\mu\left(G_{\frac{2\theta\gamma}{\eta}}^-(u, \Omega) \cap S_\phi(x_0, t_0)\right) \geq \left[(1 - \epsilon)^{\frac{1}{n}} - \frac{\eta}{2n\theta\gamma} \left(\int_{S_\phi(x_0, t_0)} \left| \frac{f}{\det D^2\phi} \right|^n d\mu \right)^{\frac{1}{n}} \right]^n \mu(S_\phi(x_0, t_0))$$

for all $\eta \leq \eta(\epsilon, n, \lambda, \Lambda)$. Here $\mu := M\phi$ and $\theta = \theta(n, \lambda, \Lambda) > 1$ is the engulfing constant given by [G, Theorem 3.3.7].

Proof. Let T normalize $S_\phi(x_0, t_0)$, and for $y \in T(\Omega)$ we set

$$\tilde{\phi}(y) = \frac{1}{t_0} \left[\phi(T^{-1}y) - \phi(x_0) - \nabla\phi(x_0) \cdot (T^{-1}y - x_0) - t_0 \right] \quad \text{and} \quad \tilde{u}(y) = \frac{1}{2\theta t_0} u(T^{-1}y).$$

It follows that $\tilde{\Omega} := T(S_\phi(x_0, t_0))$ is normalized, $\tilde{\phi} = 0$ on $\partial\tilde{\Omega}$ and $\lambda' \leq \det D^2\tilde{\phi} \leq \Lambda'$ in $\tilde{\Omega}$, where λ' and Λ' depend only on n, λ and Λ .

Let $\bar{x} \in S_\phi(x_0, t_0) \cap G_\gamma(u, \Omega)$ be such that $S_\phi(\bar{x}, \theta t_0) \Subset U$, and define $\bar{y} = T\bar{x}$. Then $-\gamma d(x, \bar{x})^2 \leq u(x) - u(\bar{x}) - \nabla u(\bar{x}) \cdot (x - \bar{x}) \leq \gamma d(x, \bar{x})^2$ for all x in Ω . Hence by changing variables we get

$$(3.18) \quad -\gamma \frac{d(T^{-1}y, T^{-1}\bar{y})^2}{2\theta t_0} \leq \tilde{u}(y) - \tilde{u}(\bar{y}) - \nabla\tilde{u}(\bar{y}) \cdot (y - \bar{y}) \leq \gamma \frac{d(T^{-1}y, T^{-1}\bar{y})^2}{2\theta t_0}, \quad \forall y \in T(\Omega).$$

Since $\bar{x} \in S_\phi(x_0, t_0)$, we have $S_\phi(x_0, t_0) \subset S_\phi(\bar{x}, \theta t_0)$ by the engulfing property. So, if $x \in S_\phi(x_0, t_0)$, then $d(x, \bar{x})^2 \leq \theta t_0$, and consequently $d(T^{-1}y, T^{-1}\bar{y})^2 \leq \theta t_0$ for all $y \in T(S_\phi(x_0, t_0))$. This together with (3.18) gives

$$-\frac{\gamma}{2} \leq \tilde{u}(y) - \tilde{u}(\bar{y}) - \nabla\tilde{u}(\bar{y}) \cdot (y - \bar{y}) \leq \frac{\gamma}{2} \quad \text{in } \tilde{\Omega}.$$

Hence if $v(y) := \frac{1}{\gamma} \left[\tilde{u}(y) - \tilde{u}(\bar{y}) - \nabla\tilde{u}(\bar{y}) \cdot (y - \bar{y}) + \frac{\gamma}{2} \right]$ for $y \in T(\Omega)$, then $0 \leq v \leq 1$ in $\tilde{\Omega}$.

Let $0 < \alpha < 1$. There exists $\eta(\alpha) = \eta(\alpha, n, \lambda, \Lambda) > 0$ such that if $\bar{y} \in \tilde{\Omega}_\alpha$, then

$$(3.19) \quad \tilde{\phi}(\bar{y}) + \nabla\tilde{\phi}(\bar{y}) \cdot (y - \bar{y}) + \eta(\alpha) \left(\theta + \frac{1}{2} \right) < 0, \quad \text{for all } y \in \overline{\tilde{\Omega}}.$$

Define $w_\alpha(y) = \eta(\alpha)v(y) + \tilde{\phi}(y)$. Let γ_α be the convex envelope of w_α in $\tilde{\Omega}$, and

$$C_\alpha = \left\{ \tilde{y} \in \tilde{\Omega} : w_\alpha(\tilde{y}) = \gamma_\alpha(\tilde{y}), \text{ and } \exists \ell \text{ supporting hyperplane to } \gamma_\alpha \text{ at } \tilde{y}, \right. \\ \left. \text{with } \ell < -\eta(\alpha) \left(\theta - \frac{1}{2} \right) \text{ in } \tilde{\Omega} \right\}.$$

Claim 1. $\nabla\tilde{\phi}(\tilde{\Omega}_\alpha) = \nabla(\tilde{\phi} + \eta(\alpha))(C_\alpha) \subset \nabla w_\alpha(C_\alpha)$.

The proof of this is similar to that of Claim 1 in Lemma 3.2.

Claim 2. $C_\alpha \subset T\left(G_{2\theta\gamma/\eta(\alpha)}^-(u, \Omega) \cap S_\phi(x_0, t_0)\right)$ for every $0 < \alpha < 1$.

Proof of Claim 2. Let $\tilde{y} \in C_\alpha$. There exists ℓ affine such that $\eta(\alpha)v(y) + \tilde{\phi}(y) \geq \ell(y)$ for all $y \in T(S_\phi(x_0, t_0))$, and with equality at $y = \tilde{y}$, and $\ell < -\eta(\alpha)(\theta - 1/2)$

in $T(S_\phi(x_0, t_0))$. Since $\tilde{\phi}(y) = \tilde{\ell}_{\tilde{y}}(y) + \frac{1}{t_0} d(T^{-1}y, T^{-1}\tilde{y})^2 \forall y \in T(\Omega)$ where $\tilde{\ell}_{\tilde{y}}(y) := \tilde{\phi}(\tilde{y}) + \nabla\tilde{\phi}(\tilde{y}) \cdot (y - \tilde{y})$, we then have

$$(3.20) \quad \eta(\alpha)v(y) \geq \ell(y) - \tilde{\ell}_{\tilde{y}}(y) - \frac{1}{t_0} d(T^{-1}y, T^{-1}\tilde{y})^2 =: g(y) \quad \forall y \in T(S_\phi(x_0, t_0))$$

with equality at $y = \tilde{y}$. Our goal is to extend (3.20) to the set $T(\Omega)$. We claim that

$$(3.21) \quad g(y) < \frac{\eta(\alpha)}{2} \left(1 - \frac{d(T^{-1}y, T^{-1}\tilde{y})^2}{t_0\theta} \right) \quad \forall y \in T(\Omega) \setminus T(S_\phi(x_0, t_0)).$$

Assume this claim for a moment. Notice that from (3.18) we have that

$$(3.22) \quad \eta(\alpha)v(y) \geq \frac{\eta(\alpha)}{2} \left(1 - \frac{d(T^{-1}y, T^{-1}\tilde{y})^2}{t_0\theta} \right) \quad \forall y \in T(\Omega),$$

and therefore (3.20) holds for all $y \in T(\Omega)$. Using (3.20), the fact $g(y) = \eta(\alpha)v(\tilde{y}) + \eta(\alpha)\nabla v(\tilde{y}) \cdot (y - \tilde{y}) - \frac{1}{t_0} d(T^{-1}y, T^{-1}\tilde{y})^2$ and the definition of v , we obtain

$$\tilde{u}(y) \geq \tilde{u}(\tilde{y}) + \nabla\tilde{u}(\tilde{y}) \cdot (y - \tilde{y}) - \frac{\gamma}{t_0\eta(\alpha)} d(T^{-1}y, T^{-1}\tilde{y})^2 \quad \forall y \in T(\Omega).$$

Thus we have shown that

$$\begin{aligned} C_\alpha &\subset \{ \tilde{y} \in T(S_\phi(x_0, t_0)) : \tilde{u}(y) \geq \tilde{u}(\tilde{y}) + \nabla\tilde{u}(\tilde{y}) \cdot (y - \tilde{y}) - \frac{\gamma}{t_0\eta(\alpha)} d(T^{-1}y, T^{-1}\tilde{y})^2 \forall y \in T(\Omega) \} \\ &= T\left\{ \tilde{x} \in S_\phi(x_0, t_0) : u(x) \geq u(\tilde{x}) + \nabla u(\tilde{x}) \cdot (x - \tilde{x}) - \frac{2\theta\gamma}{\eta(\alpha)} d(x, \tilde{x})^2 \quad \forall x \in \Omega \right\} \\ &= T\left(G_{\frac{2\theta\gamma}{\eta(\alpha)}}^-(u, \Omega) \cap S_\phi(x_0, t_0) \right). \end{aligned}$$

So Claim 2 holds as long as (3.21) is proved. Observe that (3.21) is equivalent to

$$(3.23) \quad B := \left\{ y \in T(\Omega) : g(y) \geq \frac{\eta(\alpha)}{2} \left(1 - \frac{d(T^{-1}y, T^{-1}\tilde{y})^2}{t_0\theta} \right) \right\} \subset T(S_\phi(x_0, t_0)).$$

Since $\eta(\alpha)/2\theta < 1$, we have that the function

$$\begin{aligned} &-g(y) + \frac{\eta(\alpha)}{2} \left(1 - \frac{d(T^{-1}y, T^{-1}\tilde{y})^2}{t_0\theta} \right) \\ &= -\ell(y) + \tilde{\ell}_{\tilde{y}}(y) + \frac{1}{t_0} d(T^{-1}y, T^{-1}\tilde{y})^2 + \frac{\eta(\alpha)}{2} \left(1 - \frac{d(T^{-1}y, T^{-1}\tilde{y})^2}{t_0\theta} \right) \\ &= -\ell(y) + \frac{\eta(\alpha)}{2} + \frac{\eta(\alpha)}{2\theta} [\tilde{\phi}(\tilde{y}) + \nabla\tilde{\phi}(\tilde{y}) \cdot (y - \tilde{y})] + \left(1 - \frac{\eta(\alpha)}{2\theta} \right) \tilde{\phi}(\tilde{y}) \end{aligned}$$

is convex and hence B is connected. Moreover, $\tilde{y} \in B \cap T(S_\phi(x_0, t_0))$ by (3.22) and since $g(\tilde{y}) = \eta(\alpha)v(\tilde{y})$. Thus, (3.23) will follow if

$$(3.24) \quad B \cap \partial T(S_\phi(x_0, t_0)) = \emptyset.$$

Recall that $\ell < -\eta(\alpha)(\theta - 1/2)$ in $T(S_\phi(x_0, t_0))$, and $\tilde{\ell}_{\tilde{y}}(y) + \frac{1}{t_0} d(T^{-1}y, T^{-1}\tilde{y})^2 = \tilde{\phi}(y) = 0$ on $\partial T(S_\phi(x_0, t_0))$. In addition, $d(T^{-1}y, T^{-1}\tilde{y})^2 \leq \theta t_0$ in $T(S_\phi(x_0, t_0))$ since $S_\phi(x_0, t_0) \subset S_\phi(\bar{x}, \theta t_0)$. Therefore, if $y \in \partial T(S_\phi(x_0, t_0))$ then

$$\begin{aligned} & -g(y) + \frac{\eta(\alpha)}{2} \left(1 - \frac{d(T^{-1}y, T^{-1}\tilde{y})^2}{t_0\theta} \right) \\ & = -\ell(y) + \tilde{\ell}_{\tilde{y}}(y) + \frac{1}{t_0} d(T^{-1}y, T^{-1}\tilde{y})^2 + \frac{\eta(\alpha)}{2} \left(1 - \frac{d(T^{-1}y, T^{-1}\tilde{y})^2}{t_0\theta} \right) \geq \eta(\alpha) \left(\theta - \frac{1}{2} \right) > 0, \end{aligned}$$

and hence (3.24) holds as desired. This completes the proof of (3.21), and so Claim 2 is proved.

The lemma now follows by applying Lemma 3.1 with $\Omega \rightsquigarrow \tilde{\Omega}$, $u \rightsquigarrow \eta(\alpha)v$, $\phi \rightsquigarrow \tilde{\phi}$, $E = C_\alpha$ and using Claim 1 and Claim 2. The detailed calculations are the same as those in Lemma 3.2. \square

3.2. Initial power decay for the linearized Monge-Ampère equation. We next use Lemma 3.2 and Lemma 3.3 to derive a small power decay estimate. To achieve this, the covering result proved in [CG2] is essential.

Proposition 3.4. *Let U be a normalized convex domain and Ω be a bounded convex set such that $U \subset \Omega$. Let $\phi \in C^1(\Omega) \cap W_{loc}^{2,n}(U)$ be a convex function satisfying $\lambda \leq \det D^2\phi \leq \Lambda$ in U and $\phi = 0$ on ∂U . Suppose $u \in C(\Omega) \cap W_{loc}^{2,n}(U) \cap C^1(U)$, $|u| \leq 1$ in Ω and $\mathcal{L}_\phi u = f$ in U with $\|f/\det D^2\phi\|_{L^\infty(U,\mu)} \leq 1$. Then for any $0 < \alpha < 1$, there exist $C, \tau > 0$ depending only on α, n, λ and Λ such that*

$$\mu(U_\alpha \setminus G_\beta(u, \Omega)) \leq \frac{C}{\beta^\tau} \quad \text{for all } \beta \text{ large,}$$

where U_α is defined as in (2.6).

Proof. Let $0 < \epsilon < 1/2$ and $\eta(\epsilon, n, \lambda, \Lambda)$ be the smallest of the constants in Lemma 3.2 and Lemma 3.3. Next fix $0 < \eta \leq \eta(\epsilon, n, \lambda, \Lambda)$ small so that $[(1 - \epsilon)^{1/n} - C\eta]^n \geq 1 - 2\epsilon$, where $C = C(n, \lambda, \Lambda)$. Applying Lemma 3.2 to the functions $\frac{u+1}{2}$ and $\frac{-u+1}{2}$, and noticing that $G_N^-(\frac{u+1}{2}, \Omega) = G_{2N}^-(u, \Omega)$ and $G_N^-(\frac{-u+1}{2}, \Omega) = G_{2N}^+(u, \Omega)$, we obtain

$$\begin{aligned} \mu(S_\phi(x_0, t_0) \cap G_{2/\eta t_0}^-(u, \Omega)) & \geq [(1 - \epsilon)^{1/n} - C\eta]^n \mu(S_\phi(x_0, t_0)) \geq (1 - 2\epsilon) \mu(S_\phi(x_0, t_0)), \\ \mu(S_\phi(x_0, t_0) \cap G_{2/\eta t_0}^+(u, \Omega)) & \geq [(1 - \epsilon)^{1/n} - C\eta]^n \mu(S_\phi(x_0, t_0)) \geq (1 - 2\epsilon) \mu(S_\phi(x_0, t_0)) \end{aligned}$$

for any $S_\phi(x_0, t_0) \Subset U$. Taking $M := 2\theta/\eta$, it then follows that

$$\begin{aligned} \mu(S_\phi(x_0, t_0) \setminus G_{M/\theta t_0}(u, \Omega)) & \leq \mu(S_\phi(x_0, t_0) \setminus G_{M/\theta t_0}^+(u, \Omega)) + \mu(S_\phi(x_0, t_0) \setminus G_{M/\theta t_0}^-(u, \Omega)) \\ & \leq 4\epsilon \mu(S_\phi(x_0, t_0)) \end{aligned}$$

as long as $S_\phi(x_0, t_0) \Subset U$.

Set $\alpha_0 := \frac{\alpha+1}{2}$. Assume $\alpha_2 < \alpha_1 < \alpha_0$ are such that there exist $\eta_2 < \eta_1$ with the property: if $x \in U_{\alpha_2}$ and $t \leq \eta_2$ then $S_\phi(x, t) \subset U_{\alpha_1}$; and if $x \in U_{\alpha_1}$ and $t \leq \eta_1$ then $S_\phi(x, t) \subset U_{\alpha_0}$. Let $h \geq 1/\eta_0$ satisfy $1/\theta h \leq \eta_2$. For $x_0 \in U_{\alpha_2} \setminus G_{hM}(u, \Omega)$, define

$$g(t) := \frac{\mu\left((U_{\alpha_2} \setminus G_{hM}(u, \Omega)) \cap S_\phi(x_0, t)\right)}{\mu(S_\phi(x_0, t))}, \quad t > 0.$$

We have $\lim_{t \rightarrow 0} g(t) = 1$. Also, if $1/\theta h \leq t < \eta_1$, then $S_\phi(x_0, t) \subset U_{\alpha_0}$ and

$$\begin{aligned} \mu\left((U_{\alpha_2} \setminus G_{hM}(u, \Omega)) \cap S_\phi(x_0, t)\right) &\leq \mu(S_\phi(x_0, t) \setminus G_{hM}(u, \Omega)) \\ &\leq \mu(S_\phi(x_0, t) \setminus G_{M/\theta t}(u, \Omega)) \leq 4\epsilon \mu(S_\phi(x_0, t)), \end{aligned}$$

since $G_{M/\theta t}(u, \Omega) \subset G_{hM}(u, \Omega)$. Therefore $g(t) \leq 4\epsilon$ for $t \in [1/\theta h, \eta_1)$ and so by continuity of g , there exists $t_{x_0} \leq 1/\theta h$ satisfying $g(t_{x_0}) = 4\epsilon$. Thus, we have shown that for any $x_0 \in U_{\alpha_2} \setminus G_{hM}(u, \Omega)$ there is $t_{x_0} \leq 1/\theta h$ such that

$$(3.25) \quad \mu\left((U_{\alpha_2} \setminus G_{hM}(u, \Omega)) \cap S_\phi(x_0, t_{x_0})\right) = 4\epsilon \mu(S_\phi(x_0, t_{x_0})).$$

We now claim that (3.25) implies

$$(3.26) \quad S_\phi(x_0, t_{x_0}) \subset \left(U_{\alpha_1} \setminus G_h(u, \Omega)\right) \cup \left\{x \in U_{\alpha_0} : \mathcal{M}_\mu((f/\det D^2\phi)^n)(x) > (2\theta nh)^n\right\}.$$

Otherwise, and since $x_0 \in U_{\alpha_2}$ and $t_{x_0} \leq 1/\theta h \leq \eta_2$, we have $S_\phi(x_0, t_{x_0}) \subset U_{\alpha_1}$ and there exists $\bar{x} \in S_\phi(x_0, t_{x_0}) \cap G_h(u, \Omega)$ such that $\mathcal{M}_\mu((f/\det D^2\phi)^n)(\bar{x}) \leq (2\theta nh)^n$. Note also that $S_\phi(\bar{x}, \theta t_{x_0}) \Subset U$ as $\bar{x} \in U_{\alpha_0}$ and $\theta t_{x_0} \leq 1/h \leq \eta_0$. Then by Lemma 3.3 applied to u and $-u$ and by our choice of η , we obtain

$$\begin{aligned} (1 - 2\epsilon) \mu(S_\phi(x_0, t_{x_0})) &< \mu(S_\phi(x_0, t_{x_0}) \cap G_{hM}^-(u, \Omega)), \\ (1 - 2\epsilon) \mu(S_\phi(x_0, t_{x_0})) &< \mu(S_\phi(x_0, t_{x_0}) \cap G_{hM}^+(u, \Omega)). \end{aligned}$$

Hence

$$\mu\left((U_{\alpha_2} \setminus G_{hM}(u, \Omega)) \cap S_\phi(x_0, t_{x_0})\right) \leq \mu\left(S_\phi(x_0, t_{x_0}) \setminus G_{hM}(u, \Omega)\right) < 4\epsilon \mu(S_\phi(x_0, t_{x_0})),$$

a contradiction with (3.25). So (3.26) is proved and we can apply the covering result [G, Theorem 6.3.3] to conclude that

$$(3.27) \quad \begin{aligned} &\mu(U_{\alpha_2} \setminus G_{hM}(u, \Omega)) \\ &\leq 2\sqrt{\epsilon} \left[\mu(U_{\alpha_1} \setminus G_h(u, \Omega)) + \mu\left\{x \in U_{\alpha_0} : \mathcal{M}_\mu((f/\det D^2\phi)^n)(x) > (2\theta nh)^n\right\} \right], \end{aligned}$$

as long as $\alpha_2 < \alpha_1 < \alpha_0$ are such that $\eta_2 < \eta_1$, and $h \geq 1/\eta_0$ satisfy $1/\theta h \leq \eta_2$.

For $k \in \mathbb{N}$, set

$$a_k := \mu(U_{\alpha_k} \setminus G_{M^k}(u, \Omega)) \quad \text{and} \quad b_k := \mu\left\{x \in U_{\alpha_0} : \mathcal{M}_\mu((f/\det D^2\phi)^n)(x) > (2\theta nM^k)^n\right\},$$

where α_k will be defined inductively in the sequel. First fix α_1 so that $2\alpha_0 - 1 < \alpha_1 < \alpha_0$ and take $\eta_1 := C_0(\alpha_0 - \alpha_1)^\gamma$, where C_0 and γ are the constants in Lemma 2.1.

Let $h = M$, and set $\alpha_2 = \alpha_1 - \frac{1}{(C_0\theta M)^{1/\gamma}}$. Then $\frac{1}{\theta h} = \frac{1}{\theta M} = C_0(\alpha_1 - \alpha_2)^\gamma =: \eta_2$, and so from Lemma 2.1 and (3.27) we get

$$a_2 \leq 2\sqrt{\epsilon}(a_1 + b_1).$$

Next let $h = M^2$ and $\alpha_3 = \alpha_2 - \frac{1}{(C_0\theta M^2)^{1/\gamma}}$, so $\frac{1}{\theta h} = C_0(\alpha_2 - \alpha_3)^\gamma =: \eta_3$. Then

$$a_3 \leq 2\sqrt{\epsilon}(a_2 + b_2).$$

Continuing in this way we let $h = M^k$ and $\alpha_{k+1} = \alpha_k - \frac{1}{(C_0\theta M^k)^{1/\gamma}}$. Then $\frac{1}{\theta h} = C_0(\alpha_k - \alpha_{k+1})^\gamma =: \eta_k$, and $a_{k+1} \leq 2\sqrt{\epsilon}(a_k + b_k)$. These imply that

$$a_{k+1} \leq (2\sqrt{\epsilon})^k a_1 + \sum_{i=1}^k (2\sqrt{\epsilon})^{(k+1)-i} b_i.$$

On the other hand, $\alpha_{k+1} = \alpha_1 - \sum_{j=1}^k \frac{1}{(C_0\theta M^j)^{1/\gamma}} \geq \alpha_1 - \frac{1}{(C_0\theta)^{1/\gamma}} \frac{1}{M^{1/\gamma-1}} \geq 2\alpha_0 - 1$ by choosing η even smaller depending on α (recall that $M = 2\theta/\eta$). Therefore, we obtain

$$\mu(U_{2\alpha_0-1} \setminus G_{M^{k+1}}(u, \Omega)) \leq \mu(U_{\alpha_{k+1}} \setminus G_{M^{k+1}}(u, \Omega)) \leq (2\sqrt{\epsilon})^k a_1 + \sum_{i=1}^k (2\sqrt{\epsilon})^{(k+1)-i} b_i$$

for all $k = 1, 2, \dots$. Moreover,

$$b_i \leq \frac{C(n, \lambda, \Lambda)}{M^{ni}} \int_U \left| \frac{f}{\det D^2 \phi} \right|^n d\mu(x) \leq C(n, \lambda, \Lambda) M^{-ni}$$

because \mathcal{M}_μ is of weak type 1 – 1 (see [GT, Theorem 2.9]). Thus, by setting $m_0 := \max\{2\sqrt{\epsilon}, M^{-n}\}$ we then have

$$\mu(U_{2\alpha_0-1} \setminus G_{M^{k+1}}(u, \Omega)) \leq m_0^k a_1 + C m_0^{k+1} \leq C(\epsilon, n, \lambda, \Lambda) m_0^{k+1} (1 + k).$$

Writing $m_1 = \sqrt{m_0}$ and since $m_0 < 1$, we conclude that $m_0^{k+1} (1 + k) \leq C'(m_0) m_1^{k+1}$ and so $\mu(U_{2\alpha_0-1} \setminus G_{M^{k+1}}(u, \Omega)) \leq C m_1^{k+1}$. Now for any $\beta \geq M^2$, pick $k \in \mathbb{N}$ such that $M^{k+1} \leq \beta < M^{k+2}$, then $k + 1 \leq \log_M \beta < k + 2$ and

$$\begin{aligned} \mu(U_\alpha \setminus G_\beta(u, \Omega)) &= \mu(U_{2\alpha_0-1} \setminus G_\beta(u, \Omega)) \\ &\leq \mu(U_{2\alpha_0-1} \setminus G_{M^{k+1}}(u, \Omega)) \leq C m_1^{k+1} \leq \frac{C}{m_1} \beta^{\log_M m_1}. \end{aligned}$$

□

4. L^p ESTIMATES FOR SECOND DERIVATIVES

We established in Proposition 3.4 that

$$\mu(U_\alpha \setminus G_\beta(u, \Omega)) \leq C\beta^{-\tau}$$

when $\lambda \leq \det D^2 \phi \leq \Lambda$. This power decay estimate is very poor as $\tau > 0$ is small. However, we will demonstrate in this section that τ can be taken to be any finite number provided that $\det D^2 \phi$ is sufficiently close to the constant 1 in L^∞ norm.

In order to perform this acceleration process, the following approximation lemma is crucial. This lemma is a variant of [GN, Lemma 4.1] and allows us to compare explicitly two solutions originating from two different linearized Monge-Ampère equations. We assume below that $\phi, w \in C(\bar{U})$ are convex functions satisfying $\frac{1}{2} \leq \det D^2\phi \leq \frac{3}{2}$, $\det D^2w = 1$ in U and $\phi = w = 0$ on ∂U . Also the matrices of cofactors of $D^2\phi$ and D^2w are denoted by Φ and W respectively.

Lemma 4.1. *Let U be a normalized convex domain and $u \in W_{loc}^{2,n}(U) \cap C(\bar{U})$ be a solution of $\Phi_{ij}D_{ij}u = f$ in U with $|u| \leq 1$ in U . Assume $0 < \alpha_1 < 1$ and $h \in W_{loc}^{2,n}(U_{\alpha_1}) \cap C(\bar{U}_{\alpha_1})$ is a solution of*

$$\begin{cases} W_{ij}D_{ij}h = 0 & \text{in } U_{\alpha_1} \\ h = u & \text{on } \partial U_{\alpha_1}. \end{cases}$$

Then there exists $\gamma \in (0, 1)$ depending only on n such that for any $0 < \alpha_2 < \alpha_1$, we have

$$\|u - h\|_{L^\infty(U_{\alpha_2})} + \|f - \text{trace}([\Phi - W]D^2h)\|_{L^n(U_{\alpha_2})} \leq C(\alpha_1, \alpha_2, n) \left\{ \|\Phi - W\|_{L^n(U_{\alpha_1})}^\gamma + \|f\|_{L^n(U)} \right\}$$

provided that $\|\Phi - W\|_{L^n(U_{\alpha_1})} \leq (\alpha_1 - \alpha_2)^{\frac{2n}{1+(n-1)\gamma}}$.

Proof. Let $0 < \alpha < \alpha_1$. We first claim that

$$(4.28) \quad \delta_1 := c_n(\alpha_1 - \alpha)^n \leq \text{dist}(x, \partial U_{\alpha_1}) \leq 2n \min \{1, \alpha^{-1}(\alpha_1 - \alpha)\} =: \delta_2 \quad \forall x \in \partial U_\alpha.$$

To prove (4.28), let x_0 be the minimum point of ϕ in U . Then $U_\alpha = S_\phi(x_0, -\alpha\phi(x_0))$, $U_{\alpha_1} = S_\phi(x_0, -\alpha_1\phi(x_0))$, and $C_1(n) \leq |\phi(x_0)| \leq C_2(n)$ by [G, Proposition 3.2.3]. For any $x \in \partial U_\alpha$, by applying Aleksandrov's estimate (see [G, Theorem 1.4.2]) to the function $\tilde{\phi} := \phi - (1 - \alpha_1)\phi(x_0)$ we get $\text{dist}(x, \partial U_{\alpha_1})^{1/n} \geq C_n|\tilde{\phi}(x)| = C_n(\alpha - \alpha_1)\phi(x_0)$ yielding the first inequality in (4.28). For the second inequality, let $x \in \partial U_\alpha$ and choose y be such that $x = (1 - \frac{\alpha}{\alpha_1})x_0 + \frac{\alpha}{\alpha_1}y$. Then $y \notin U_{\alpha_1}$ since whenever $y \in U$ we have $(1 - \alpha_1)\phi(x_0) \leq \phi(y)$ as $(1 - \alpha)\phi(x_0) = \phi(x) \leq (1 - \frac{\alpha}{\alpha_1})\phi(x_0) + \frac{\alpha}{\alpha_1}\phi(y)$ by the convexity of ϕ . Therefore, we infer that $\text{dist}(x, \partial U_{\alpha_1}) \leq |y - x| = (\frac{\alpha_1}{\alpha} - 1)|x - x_0| \leq 2n(\frac{\alpha_1}{\alpha} - 1)$ which gives the desired result.

By Caffarelli-Gutiérrez interior Hölder estimates (see [GN, estimate (2.2) and Corollary 2.6]) there exists $\beta \in (0, 1)$ depending only on n such that

$$(4.29) \quad \|u\|_{C^\beta(\bar{U}_{\alpha_1})} \leq C(\alpha_1, n)(1 + \|f\|_{L^n(U)}).$$

Next notice that Pogorelov's estimates imply that $\lambda(\alpha_1, n)I \leq W \leq \Lambda(\alpha_1, n)I$ in U_{α_1} . Therefore, by using standard boundary Hölder estimates for linear uniformly elliptic equations (see [GiT, Corollary 9.29] and [CC, Proposition 4.13]) and (4.29), we obtain

$$(4.30) \quad \|h\|_{C^{\beta/2}(\bar{U}_{\alpha_1})} \leq C'(\alpha_1, n)\|u\|_{C^\beta(\partial U_{\alpha_1})} \leq C(\alpha_1, n)(1 + \|f\|_{L^n(U)}).$$

Now for any $x \in \partial U_\alpha$, by (4.28) we can take $y \in \partial U_{\alpha_1}$ such that $|x - y| \leq \delta_2$. Then since $u - h = 0$ on ∂U_{α_1} , we get from (4.29) and (4.30) that

$$\begin{aligned} |(u - h)(x)| &= |(u - h)(x) - (u - h)(y)| \leq |u(x) - u(y)| + |h(x) - h(y)| \\ &\leq C(\alpha_1, n) \delta_2^{\beta/2} (1 + \|f\|_{L^n(U)}). \end{aligned}$$

That is,

$$(4.31) \quad \|u - h\|_{L^\infty(\partial U_\alpha)} \leq C(\alpha_1, n) \delta_2^{\beta/2} (1 + \|f\|_{L^n(U)}).$$

We claim that

$$(4.32) \quad \|D^2 h\|_{L^\infty(U_\alpha)} \leq C(\alpha_1, n) \delta_1^{\frac{\beta}{2}-2} (1 + \|f\|_{L^n(U)}).$$

Indeed, let $x_0 \in U_\alpha$ be arbitrary and take $x_1 \in \partial B_{\delta_1/2}(x_0)$. Since $B_{\delta_1/2}(x_0) \Subset U_{\alpha_1}$ by (4.28) and $W_{ij} D_{ij}(h - h(x_1)) = W_{ij} D_{ij} h = 0$ in U_{α_1} , we can apply interior C^2 -estimates (see [GN, Theorem 2.7]) to $h - h(x_1)$ in $B_{\delta_1/2}(x_0)$ and obtain

$$\|D^2 h(x_0)\| \leq C'(\alpha_1, n) \delta_1^{-2} \sup_{B_{\delta_1/2}(x_0)} |h - h(x_1)| \leq C(\alpha_1, n) \delta_1^{-2} \delta_1^{\beta/2} (1 + \|f\|_{L^n(U)})$$

giving (4.32).

Observe that $u - h \in W_{loc}^{2,n}(U)$ is a solution of

$$\Phi_{ij} D_{ij}(u - h) = f - \Phi_{ij} D_{ij} h = f - [\Phi_{ij} - W_{ij}] D_{ij} h =: F \quad \text{in } U_{\alpha_1}.$$

Hence if we let $\epsilon := \|\Phi - W\|_{L^n(U_{\alpha_1})}$, then it follows from the ABP estimate (see [GN, Theorem 2.4]), (4.31) and (4.32) that

$$\begin{aligned} \|u - h\|_{L^\infty(U_\alpha)} + \|F\|_{L^n(U_\alpha)} &\leq \|u - h\|_{L^\infty(\partial U_\alpha)} + C_n \|F\|_{L^n(U_{\alpha_1})} \\ &\leq \|u - h\|_{L^\infty(\partial U_\alpha)} + C_n \|D^2 h\|_{L^\infty(U_\alpha)} \|\Phi - W\|_{L^n(U_{\alpha_1})} + C_n \|f\|_{L^n(U_{\alpha_1})} \\ &\leq C(\alpha_1, n) \left[\alpha^{-\beta/2} (\alpha_1 - \alpha)^{\beta/2} + (\alpha_1 - \alpha)^{n(\frac{\beta}{2}-2)} \epsilon \right] (1 + \|f\|_{L^n(U)}) + C_n \|f\|_{L^n(U)}. \end{aligned}$$

By taking $\alpha := \alpha_1 - \epsilon^{\frac{2}{4n-(n-1)\beta}}$, this yields

$$\|u - h\|_{L^\infty(U_\alpha)} + \|F\|_{L^n(U_\alpha)} \leq C(\alpha_1, n) (\alpha^{-\beta/2} + 1) \epsilon^\gamma (1 + \|f\|_{L^n(U)}) + C_n \|f\|_{L^n(U)}$$

with $\gamma := \frac{\beta}{4n-(n-1)\beta}$. From this we deduce the lemma as $\epsilon \leq (\alpha_1 - \alpha_2)^{\frac{2n}{1+(n-1)\gamma}}$ by the assumption. \square

4.1. Improved density estimates. In this subsection we will use Lemma 4.1 to improve the power decay of $\mu(U_\alpha \setminus G_\beta(u, \Omega))$. To this end, the next lemma plays an important role.

Lemma 4.2. *Let $0 < \epsilon < 1/2$, $0 < \alpha_0 < 1$, U be a normalized convex domain and Ω be a bounded convex set such that $U \subset \Omega$. Let $\phi \in C^1(\Omega) \cap W_{loc}^{2,n}(U)$ be a convex function satisfying $1 - \epsilon \leq \det D^2 \phi \leq 1 + \epsilon$ in U and $\phi = 0$ on ∂U . Suppose $u \in C(\Omega) \cap W_{loc}^{2,n}(U) \cap C^1(U)$ is a solution of $\mathcal{L}_\phi u = f$ in U with $|u| \leq 1$ in U and $|u(x)| \leq C^* d(x, x_0)^2$ in $\Omega \setminus U$*

for some $x_0 \in U_{\alpha_0}$. Then for any $0 < \alpha \leq \alpha_0$, there exist $C, \tau > 0$ depending only on α, α_0 and n such that

$$|G_N(u, \Omega) \cap U_\alpha| \geq \left\{1 - C(N^{-\tau} \delta_0^\tau + \epsilon)\right\} |U_\alpha|$$

for any $N \geq N_0 = N_0(\alpha, \alpha_0, C^*, n)$ and provided that $\|\Phi - W\|_{L^n(U_{\frac{\alpha_0+1}{2}})} \leq ((1-\alpha_0)/4)^{\frac{2n}{1+(n-1)\gamma}}$.

Here W, γ are from Lemma 4.1 and

$$\delta_0 := \left(\int_{U_{\frac{\alpha_0+1}{2}}} \|\Phi - W\|^n dx \right)^{\frac{1}{n}} + \left(\int_U |f|^n dx \right)^{\frac{1}{n}}.$$

Proof. Let $h \in W_{loc}^{2,n}(U_{\frac{\alpha_0+1}{2}}) \cap C(\overline{U_{\frac{\alpha_0+1}{2}}})$ be the solution of

$$\begin{cases} W_{ij} D_{ij} h = 0 & \text{in } U_{\frac{\alpha_0+1}{2}} \\ h = u & \text{on } \partial U_{\frac{\alpha_0+1}{2}}. \end{cases}$$

By the interior $C^{1,1}$ regularity of h and Lemma 4.1, we have

$$(4.33) \quad \|h\|_{C^{1,1}(U_{\frac{3\alpha_0+1}{4}})} \leq c_e(\alpha_0, n) \|u\|_{L^\infty(U_{\frac{\alpha_0+1}{2}})} \leq c_e(\alpha_0, n),$$

$$(4.34) \quad \|u - h\|_{L^\infty(U_{\frac{3\alpha_0+1}{4}})} + \|f - \text{trace}([\Phi - W]D^2 h)\|_{L^n(U_{\frac{3\alpha_0+1}{4}})} \leq C(\alpha_0, n) \delta_0 =: \delta'_0.$$

We now consider $h|_{U_{\frac{3\alpha_0+1}{4}}}$ and then extend h outside $U_{\frac{3\alpha_0+1}{4}}$ continuously such that

$$\begin{cases} h(x) = u(x) & \forall x \in \Omega \setminus U_{\frac{\alpha_0+1}{2}}, \\ \|u - h\|_{L^\infty(\Omega)} = \|u - h\|_{L^\infty(U_{(3\alpha_0+1)/4})}. \end{cases}$$

Since by the maximum principle $\|h\|_{L^\infty(U_{\frac{3\alpha_0+1}{4}})} \leq \|u\|_{L^\infty(U)} \leq 1$, we then obtain that

$$(4.35) \quad u(x) - 2 \leq h(x) \leq u(x) + 2 \quad \text{for all } x \in \Omega.$$

We claim that if $N \geq N_0 := N_0(\alpha, \alpha_0, n)$, then

$$(4.36) \quad U_\alpha \cap A_{\sigma(\alpha)} \subset G_N(h, \Omega)$$

where $\sigma(\alpha) > 0$ is the constant given by [G, Theorem 6.1.1] and

$$A_{\sigma(\alpha)} := \left\{ \bar{x} \in U : \phi(x) \geq \phi(\bar{x}) + \nabla\phi(\bar{x}) \cdot (x - \bar{x}) + \frac{\sigma(\alpha)}{2} |x - \bar{x}|^2, \quad \forall x \in U \right\}.$$

Indeed, let $\bar{x} \in U_\alpha \cap A_{\sigma(\alpha)} \subset U_{\alpha_0}$. By (4.33) we have $|h(x) - [h(\bar{x}) + \nabla h(\bar{x}) \cdot (x - \bar{x})]| \leq c_e(\alpha_0, n) |x - \bar{x}|^2$ for all $x \in U_{\frac{3\alpha_0+1}{4}}$, and since $\bar{x} \in A_{\sigma(\alpha)}$

$$(4.37) \quad d(x, \bar{x})^2 = \phi(x) - [\phi(\bar{x}) + \nabla\phi(\bar{x}) \cdot (x - \bar{x})] \geq \frac{\sigma(\alpha)}{2} |x - \bar{x}|^2 \quad \forall x \in \bar{U}.$$

Therefore

$$(4.38) \quad |h(x) - [h(\bar{x}) + \nabla h(\bar{x}) \cdot (x - \bar{x})]| \leq \frac{2c_e(\alpha_0, n)}{\sigma(\alpha)} d(x, \bar{x})^2 \quad \forall x \in U_{\frac{3\alpha_0+1}{4}}.$$

We next show that by increasing the constant on the right hand side of (4.38), that the resulting inequality holds for all x in Ω . To see this, observe that

$$(4.39) \quad \begin{aligned} d(x, x_0)^2 &= d(x, \bar{x})^2 + [\phi(\bar{x}) - \phi(x_0) - \nabla\phi(x_0) \cdot (\bar{x} - x_0)] \\ &\quad + [\nabla\phi(\bar{x}) - \nabla\phi(x_0)] \cdot (x - \bar{x}) \\ &\leq d(x, \bar{x})^2 + C(\alpha_0, n) (1 + |x - \bar{x}|) \quad \text{for all } x \in \Omega. \end{aligned}$$

Also there exists $c(\alpha, n) > 0$ such that

$$(4.40) \quad d(x, \bar{x})^2 \geq c(\alpha, n) |x - \bar{x}| \quad \forall x \in \overline{\Omega} \setminus U.$$

Notice that $\text{dist}(U_\alpha, \partial U) \geq c_n(1-\alpha)^n$ by the Aleksandrov estimate [G, Theorem 1.4.2] and [G, Proposition 3.2.3]. Thus it follows from (4.37) and the fact $\bar{x} \in U_\alpha$ that there is $c = c(\alpha, n) > 0$ so that (4.40) holds for all $x \in \partial U$. Now for $x \in \overline{\Omega} \setminus \overline{U}$ we can choose $\hat{x} \in \partial U$ and $\lambda \in (0, 1)$ satisfying $\hat{x} = \lambda x + (1 - \lambda)\bar{x}$. Then since $d(\hat{x}, \bar{x})^2 \geq c|\hat{x} - \bar{x}|$ and the function $z \mapsto d(z, \bar{x})^2$ is convex, we obtain

$$\lambda d(x, \bar{x})^2 + (1 - \lambda)d(\bar{x}, \bar{x})^2 \geq c|\lambda x + (1 - \lambda)\bar{x} - \bar{x}| = c\lambda|x - \bar{x}|$$

which gives $d(x, \bar{x})^2 \geq c|x - \bar{x}|$ and hence (4.40) is proved.

We are ready to show that (4.38) holds for all $x \in \Omega$ but with a bigger constant on the right hand side. Let $x \in \Omega \setminus U_{\frac{3\alpha_0+1}{4}}$ and consider the following cases:

Case 1: $x \in U$. Then by using (4.33), (4.35) and the assumption $|u| \leq 1$ in U , we have

$$\begin{aligned} |h(x) - [h(\bar{x}) + \nabla h(\bar{x}) \cdot (x - \bar{x})]| &\leq |h(x) - h(\bar{x})| + C(\alpha_0, n) \leq |u(x) - u(\bar{x})| + C(\alpha_0, n) \\ &\leq C(\alpha_0, n) \leq C_1(\alpha_0, n) d(x, \bar{x})^2 \end{aligned}$$

where in the last inequality we have used the fact that since $\bar{x} \in U_\alpha \subset U_{\alpha_0}$ there exists $\eta(\alpha_0) > 0$ such that $S_\phi(\bar{x}, \eta(\alpha_0)) \subset U_{\frac{3\alpha_0+1}{4}}$ (see Lemma 2.1).

Case 2: $x \in \Omega \setminus U$. Then $d(x, \bar{x})^2 \geq \eta_0$ since $S_\phi(\bar{x}, \eta_0) \Subset U$ by Lemma 2.1. This together with the assumptions, (4.33), (4.35), (4.39) and (4.40) gives

$$\begin{aligned} |h(x) - [h(\bar{x}) + \nabla h(\bar{x}) \cdot (x - \bar{x})]| &\leq |h(x) - h(\bar{x})| + C(\alpha_0, n)|x - \bar{x}| \\ &\leq |u(x)| + C(\alpha_0, n)(|x - \bar{x}| + 1) \leq C^* d(x, x_0)^2 + C(\alpha_0, n)(|x - \bar{x}| + 1) \\ &\leq C^* d(x, \bar{x})^2 + C(\alpha_0, n)(|x - \bar{x}| + 1) \leq C_2(\alpha, \alpha_0, C^*, n) d(x, \bar{x})^2. \end{aligned}$$

Therefore if we choose

$$N_0 := \max \left\{ \frac{2c_e(\alpha_0, n)}{\sigma(\alpha)}, C_1(\alpha_0, n), C_2(\alpha, \alpha_0, C^*, n) \right\},$$

then it follows from the above considerations and (4.38) that

$$|h(x) - [h(\bar{x}) + \nabla h(\bar{x}) \cdot (x - \bar{x})]| \leq N_0 d(x, \bar{x})^2 \quad \text{for all } x \in \Omega.$$

This means $\bar{x} \in G_{N_0}(h, \Omega) \subset G_N(h, \Omega)$ for all $N \geq N_0$. Thus claim (4.36) is proved.

Next let

$$u'(x) := \frac{(u - h)(x)}{\delta'_0}, \quad \text{for } x \in \Omega.$$

We infer from (4.34) and the way h was initially defined and extended that

$$\begin{aligned} \|u'\|_{L^\infty(\Omega)} &= \frac{1}{\delta'_0} \|u - h\|_{L^\infty(U_{\frac{3\alpha_0+1}{4}})} \leq 1, \\ \mathcal{L}_\phi u' &= \frac{1}{\delta'_0} [\mathcal{L}_\phi u - \mathcal{L}_\phi h] = \frac{1}{\delta'_0} [f - \text{trace}([\Phi - W]D^2h)] =: f'(x) \quad \text{in } U_{\frac{3\alpha_0+1}{4}}. \end{aligned}$$

Notice that $\|f'\|_{L^n(U_{\frac{3\alpha_0+1}{4}})} \leq 1$ by (4.34). In order to apply Proposition 3.4, let T be an invertible affine map normalizing $U_{\frac{3\alpha_0+1}{4}}$. We have $C(n) \leq |\det T| \leq C'(n)$ because $|\det T|^{-2/n} \approx |U_{\frac{3\alpha_0+1}{4}}|^{2/n} \approx \frac{3\alpha_0+1}{4} |\min_U \phi| \approx 1$. Set $\tilde{U} := T(U_{\frac{3\alpha_0+1}{4}})$, $\tilde{\Omega} := T(\Omega)$ and define

$$\tilde{\phi}(y) = |\det T|^{2/n} \left[\phi(T^{-1}y) - \left(1 - \frac{3\alpha_0+1}{4}\right) \min_U \phi \right], \quad \tilde{u}(y) = u'(T^{-1}y) \quad \text{for } y \in \tilde{\Omega}.$$

Then $1 - \epsilon \leq \det D^2 \tilde{\phi} = \det D^2 \phi(T^{-1}y) \leq 1 + \epsilon$ in \tilde{U} and $\tilde{\phi} = 0$ on $\partial \tilde{U}$. Moreover since $\tilde{\Phi}(y) = |\det T|^{-2/n} \det D^2 \phi(T^{-1}y) T[D^2 \phi(T^{-1}y)]^{-1} T^t$, we obtain

$$\text{trace}(\tilde{\Phi}(y)D^2 \tilde{u}(y)) = |\det T|^{\frac{-2}{n}} \text{trace}(\Phi(T^{-1}y)D^2 u'(T^{-1}y)) = |\det T|^{\frac{-2}{n}} f'(T^{-1}y) =: \tilde{f}(y)$$

in \tilde{U} . Thus as $\tilde{\phi} \in C^1(\tilde{\Omega}) \cap W^{2,n}(\tilde{U})$, $\tilde{u} \in C(\tilde{\Omega}) \cap W^{2,n}(\tilde{U}) \cap C^1(\tilde{U})$, $\|\tilde{u}\|_{L^\infty(\tilde{\Omega})} = \|u'\|_{L^\infty(\Omega)} \leq 1$ and $\|\tilde{f}\|_{L^n(\tilde{U})} = |\det T|^{-1/n} \|f'\|_{L^n(U_{\frac{3\alpha_0+1}{4}})} \leq |\det T|^{-1/n} \leq C_n$, we can apply Proposition 3.4 to get

$$(4.41) \quad |\tilde{U}_{\frac{4\alpha}{3\alpha_0+1}} \setminus G_{\frac{N}{\delta'_0} |\det T|^{\frac{2}{n}}}(\tilde{u}, \tilde{\Omega})| \leq C(\alpha, \alpha_0, n) \left(\frac{\delta_0}{N}\right)^\tau,$$

where $\tau > 0$ depends only on α, α_0 and n . Since $\tilde{U}_{\frac{4\alpha}{3\alpha_0+1}} = T(U_\alpha)$ and

$$\tilde{d}(Tx, T\bar{x})^2 = \tilde{\phi}(Tx) - \tilde{\phi}(T\bar{x}) - \langle \nabla \tilde{\phi}(T\bar{x}), Tx - T\bar{x} \rangle = |\det T|^{\frac{2}{n}} d(x, \bar{x})^2 \quad \forall x, \bar{x} \in \Omega,$$

we have

$$\begin{aligned} (4.42) \quad & G_\beta(\tilde{u}, \tilde{\Omega}) \cap \tilde{U}_{\frac{4\alpha}{3\alpha_0+1}} \\ &= T \left\{ \bar{x} \in U_\alpha : |\tilde{u}(y) - [\tilde{u}(T\bar{x}) + \langle \nabla \tilde{u}(T\bar{x}), y - T\bar{x} \rangle]| \leq \beta \tilde{d}(y, T\bar{x})^2 \quad \forall y \in T(\Omega) \right\} \\ &= T \left\{ \bar{x} \in U_\alpha : |u'(x) - [u'(\bar{x}) + \langle \nabla u'(\bar{x}), x - \bar{x} \rangle]| \leq \beta |\det T|^{\frac{2}{n}} d(x, \bar{x})^2 \quad \forall x \in \Omega \right\} \\ &= T \left(G_{\beta |\det T|^{\frac{2}{n}}}(u', \Omega) \cap U_\alpha \right). \end{aligned}$$

It follows from (4.41), (4.42) and the fact $T(A) \setminus T(B) = T(A \setminus B)$ that

$$|U_\alpha \setminus G_{\frac{N}{\delta'_0}}(u', \Omega)| \leq C(\alpha, \alpha_0, n) \left(\frac{\delta_0}{N}\right)^\tau.$$

As $G_{\frac{N}{\delta'_0}}(u', \Omega) = G_N(u - h, \Omega)$ and $|U_\alpha| \geq c_n \alpha^{n/2}$, we then conclude

$$|U_\alpha| - |G_N(u - h, \Omega) \cap U_\alpha| = |U_\alpha \setminus G_N(u - h, \Omega)| \leq C \left(\frac{\delta_0}{N}\right)^\tau |U_\alpha|$$

yielding

$$\begin{aligned} \left\{1 - C\left(\frac{\delta_0}{N}\right)^\tau\right\} |U_\alpha| &\leq |G_N(u - h, \Omega) \cap U_\alpha| \\ &\leq |G_N(u - h, \Omega) \cap U_\alpha \cap A_{\sigma(\alpha)}| + |U_\alpha \setminus A_{\sigma(\alpha)}| \\ &\leq |G_N(u - h, \Omega) \cap U_\alpha \cap A_{\sigma(\alpha)}| + C(\alpha, n) \epsilon |U_\alpha|, \end{aligned}$$

where the last inequality is from [G, Theorem 6.1.1]. Consequently,

$$(4.43) \quad |G_N(u - h, \Omega) \cap U_\alpha \cap A_{\sigma(\alpha)}| \geq \left\{1 - C\left[\left(\frac{\delta_0}{N}\right)^\tau + \epsilon\right]\right\} |U_\alpha|.$$

We claim that

$$(4.44) \quad G_N(u - h, \Omega) \cap U_\alpha \cap A_{\sigma(\alpha)} \subset G_{2N}(u, \Omega) \cap U_\alpha$$

which together with (4.43) gives the conclusion of the lemma. To prove the claim, let $\bar{x} \in G_N(u - h, \Omega) \cap U_\alpha \cap A_{\sigma(\alpha)}$. Then $\bar{x} \in G_N(u - h, \Omega) \cap G_N(h, \Omega)$ by (4.36). Therefore (4.44) holds because

$$\begin{aligned} &|u(x) - [u(\bar{x}) + \langle \nabla u(\bar{x}), x - \bar{x} \rangle]| \\ &\leq |(u - h)(x) - [(u - h)(\bar{x}) + \langle \nabla(u - h)(\bar{x}), x - \bar{x} \rangle]| + |h(x) - [h(\bar{x}) + \langle \nabla h(\bar{x}), x - \bar{x} \rangle]| \\ &\leq 2Nd(x, \bar{x})^2 \quad \text{for all } x \in \Omega. \end{aligned}$$

This completes the proof of the lemma. \square

By using Lemma 4.2 and a localization process, we shall prove the following.

Lemma 4.3. *Let $0 < \epsilon_0 < 1$, $0 < \alpha_0 < 1$, Ω be a normalized convex domain and $u \in W_{loc}^{2,n}(\Omega) \cap C^1(\Omega)$ be a solution of $\mathcal{L}_\phi u = f$ in Ω with $\|u\|_{L^\infty(\Omega)} \leq 1$, where $\phi \in C(\overline{\Omega})$ is a convex function satisfying $\phi = 0$ on $\partial\Omega$. There exists $\epsilon > 0$ depending only on ϵ_0 , α_0 and n such that if $1 - \epsilon \leq \det D^2\phi \leq 1 + \epsilon$ in Ω , then for any $S_\phi(x_0, \frac{t_0}{\alpha_0}) \subset \Omega_{\frac{\alpha_0+1}{2}}$ we have*

$$(4.45) \quad \left|G_{\frac{N}{t_0}}(u, \Omega) \cap S_\phi(x_0, t_0)\right| \geq \left\{1 - \epsilon_0 - C\left(\frac{t_0}{N}\right)^\tau \left(\int_{S_\phi(x_0, \frac{t_0}{\alpha_0})} |f|^n dx\right)^{\frac{\tau}{n}}\right\} |S_\phi(x_0, t_0)|$$

for every $N \geq N_0$. Here C , τ and N_0 are positive constants depending only on α_0 and n .

Proof. Observe that in fact $\phi \in C^1(\Omega)$. As ϵ will be chosen small, we also have $\phi \in W^{2,n}(\Omega_{\frac{\alpha_0+1}{2}})$ by Caffarelli $W^{2,p}$ estimates (see [C3, Theorem 1] and [G, Theorem 6.4.1]). Let T be an affine map normalizing $S_\phi(x_0, \frac{t_0}{\alpha_0})$ and let $U := T(S_\phi(x_0, \frac{t_0}{\alpha_0}))$. For each $y \in T(\Omega)$, define

$$\tilde{\phi}(y) = |\det T|^{2/n} \left[\phi(T^{-1}y) - \phi(x_0) - \nabla\phi(x_0) \cdot (T^{-1}y - x_0) - \frac{t_0}{\alpha_0} \right] \text{ and } \tilde{u}(y) = u(T^{-1}y).$$

Then $1 - \epsilon \leq \det D^2\tilde{\phi} = \det D^2\phi(T^{-1}y) \leq 1 + \epsilon$ in U and $\tilde{\phi} = 0$ on ∂U . Moreover $\text{trace}(\tilde{\Phi}(y)D^2\tilde{u}(y)) = |\det T|^{\frac{2}{n}} \text{trace}(\Phi(T^{-1}y)D^2u(T^{-1}y)) = |\det T|^{\frac{2}{n}} f(T^{-1}y) =: \tilde{f}(y)$.

Thus as $\|\tilde{u}\|_{L^\infty(T(\Omega))} = \|u\|_{L^\infty(\Omega)} \leq 1$, we obtain from Lemma 4.2 with $\alpha := \alpha_0$ that

$$|G_N(\tilde{u}, T(\Omega)) \cap U_{\alpha_0}| \geq \left\{1 - C(N^{-\tau} \delta_0^\tau + \epsilon)\right\} |U_{\alpha_0}|$$

for any $N \geq N_0 = N_0(\alpha_0, n)$ and provided that $\|\tilde{\Phi} - \tilde{W}\|_{L^n(U_{\frac{\alpha_0+1}{2}})} \leq ((1 - \alpha_0)/4)^{\frac{2n}{1+(n-1)\gamma}}$, where

$$(4.46) \quad \delta_0 := \left(\int_{U_{\frac{\alpha_0+1}{2}}} \|\tilde{\Phi} - \tilde{W}\|^n dy \right)^{\frac{\gamma}{n}} + \left(\int_U |\tilde{f}|^n dy \right)^{\frac{1}{n}},$$

γ is given by Lemma 4.1 and \tilde{W} is the cofactor matrix of $D^2\tilde{w}$ with \tilde{w} is the convex function satisfying $\det D^2\tilde{w} = 1$ in U and $\tilde{w} = 0$ on ∂U . This together with Lemma 4.4 below implies that there exists $\epsilon > 0$ sufficiently small depending only on ϵ_0, α_0 and n such that

$$\begin{aligned} |G_N(\tilde{u}, T(\Omega)) \cap U_{\alpha_0}| &\geq \left\{1 - \epsilon_0 - CN^{-\tau} \left(\int_U |\tilde{f}|^n dy \right)^{\frac{\tau}{n}}\right\} |U_{\alpha_0}| \\ &= \left\{1 - \epsilon_0 - CN^{-\tau} |\det T|^{\frac{2\tau}{n}} \left(\int_{S_\phi(x_0, \frac{t_0}{\alpha_0})} |f|^n dx \right)^{\frac{\tau}{n}}\right\} |U_{\alpha_0}| \\ &\geq \left\{1 - \epsilon_0 - C \left(\frac{t_0}{N}\right)^\tau \left(\int_{S_\phi(x_0, \frac{t_0}{\alpha_0})} |f|^n dx \right)^{\frac{\tau}{n}}\right\} |U_{\alpha_0}|. \end{aligned}$$

But since $U_{\alpha_0} = T(S_\phi(x_0, t_0))$, the same calculations leading to (4.42) yield

$$G_N(\tilde{u}, T(\Omega)) \cap U_{\alpha_0} = T(G_{N|\det T|^{\frac{2}{n}}}(u, \Omega) \cap S_\phi(x_0, t_0)) \approx T(G_{\frac{N}{t_0}}(u, \Omega) \cap S_\phi(x_0, t_0)).$$

Therefore we obtain

$$\left| T(G_{\frac{N}{t_0}}(u, \Omega) \cap S_\phi(x_0, t_0)) \right| \geq \left\{1 - \epsilon_0 - C \left(\frac{t_0}{N}\right)^\tau \left(\int_{S_\phi(x_0, \frac{t_0}{\alpha_0})} |f|^n dx \right)^{\frac{\tau}{n}}\right\} |T(S_\phi(x_0, t_0))|$$

giving (4.45) for any $N \geq N_0$. \square

In the above proof, we have used the following lemma which is a strengthened version of Lemma 3.5 in [GN]. This result is proved by using a compactness argument and [GN, Lemma 3.5].

Lemma 4.4. *Given any $0 < \epsilon_0 < 1$ and $0 < \alpha < 1$, there exists $\epsilon > 0$ depending only on ϵ_0, α and n such that if $\Omega \subset \mathbb{R}^n$ is a normalized convex domain and $\phi, w \in C(\bar{\Omega})$ are convex functions satisfying*

$$\begin{cases} 1 - \epsilon \leq \det D^2\phi \leq 1 + \epsilon & \text{in } \Omega \\ \phi = 0 & \text{on } \partial\Omega \end{cases} \quad \text{and} \quad \begin{cases} \det D^2w = 1 & \text{in } \Omega \\ w = 0 & \text{on } \partial\Omega, \end{cases}$$

then

$$\|\Phi - W\|_{L^n(\Omega_\alpha)} \leq \epsilon_0,$$

where $\Omega_\alpha := \{x \in \Omega : \phi(x) < (1 - \alpha) \min_\Omega \phi\}$.

Proof. Suppose by contradiction that it is not true. Then there exist $\epsilon_0, \alpha \in (0, 1)$, $n \in \mathbb{N}$, a sequence of normalized convex domains Ω^k and sequences of convex functions $\phi_k, w_k \in C(\overline{\Omega^k})$ with

$$\begin{cases} 1 - \frac{1}{k} \leq \det D^2 \phi_k \leq 1 + \frac{1}{k} & \text{in } \Omega^k \\ \phi_k = 0 & \text{on } \partial \Omega^k \end{cases} \quad \text{and} \quad \begin{cases} \det D^2 w_k = 1 & \text{in } \Omega^k \\ w_k = 0 & \text{on } \partial \Omega^k \end{cases}$$

such that

$$(4.47) \quad \|\Phi_k - W_k\|_{L^n(\Omega_\alpha^k)} \geq \epsilon_0 \quad \text{for all } k.$$

By Blaschke selection theorem, there is a subsequence of Ω^k , still denoted by Ω^k , such that Ω^k converges in the Hausdorff metric to a normalized convex domain Ω . Also by [G, Lemma 5.3.1] we have up to a subsequence $\phi_k \rightarrow \phi$ and $w_k \rightarrow w$ uniformly on compact subsets of Ω , where $\phi, w \in C(\overline{\Omega})$ are both convex solutions to the equation

$$\begin{cases} \det D^2 w = 1 & \text{in } \Omega, \\ w = 0 & \text{on } \partial \Omega. \end{cases}$$

Thus $\phi \equiv w$ by the uniqueness of convex solutions to the Monge-Ampère equation.

Next observe that the Aleksandrov estimate [G, Theorem 1.4.2] and [G, Proposition 3.2.3] yield

$$(4.48) \quad \text{dist}(\Omega_\alpha^k, \partial \Omega^k) \geq c_n(1 - \alpha)^n =: \tau \quad \forall k.$$

For $E \subset \mathbb{R}^n$, let $E(r) := \{x \in E : \text{dist}(x, \partial E) > r\}$ and $\delta_r(E) := \{x \in \mathbb{R}^n : \text{dist}(x, E) < r\}$. We then claim that

$$(4.49) \quad \Omega_\alpha^k \subset \Omega(\tau/2) \subset \Omega(\tau/4) \subset \Omega^k \quad \text{for all } k \text{ sufficiently large.}$$

Indeed, it follows from (4.48) that $\Omega_\alpha^k \subset \Omega^k(\tau)$. Moreover since $\Omega^k \rightarrow \Omega$ in the Hausdorff metric, we have $\Omega^k \subset \delta_{\frac{\tau}{2}}(\Omega)$ for all k large (see [Sc] for the definition of the Hausdorff distance). Therefore, $\Omega_\alpha^k \subset \delta_{\frac{\tau}{2}}(\Omega)(\tau) = \Omega(\tau/2)$ giving the first inclusion in (4.49). We also infer from the Hausdorff convergence of Ω^k to Ω that $\Omega \subset \delta_{\frac{\tau}{4}}(\Omega^k)$ for all k large. This implies $\Omega(\tau/4) \subset \delta_{\frac{\tau}{4}}(\Omega^k)(\tau/4) = \Omega^k$ and the last inclusion in (4.49) is proved.

By (4.49) and [GN, Lemma 3.5] we get $\Phi_k \rightarrow \Phi$ in $L^n(\Omega(\tau/2))$ and $W_k \rightarrow W$ in $L^n(\Omega(\tau/2))$, where Φ is the cofactor matrix of $D^2 \phi$ and W is the cofactor matrix of $D^2 w$. Since $\Phi \equiv W$, this yields $\Phi_k - W_k \rightarrow 0$ in $L^n(\Omega(\tau/2))$. Combining this with the first inclusion in (4.49) we obtain

$$\lim_{k \rightarrow \infty} \|\Phi_k - W_k\|_{L^n(\Omega_\alpha^k)} = 0,$$

which is a contradiction with (4.47) and the proof is complete. \square

In the next lemma, we no longer require $\|u\|_{L^\infty(\Omega)} \leq 1$ as in Lemma 4.3.

Lemma 4.5. *Let $0 < \epsilon_0 < 1$, $0 < \alpha_0 < 1$, Ω be a normalized convex domain and $u \in W_{loc}^{2,n}(\Omega) \cap C^1(\Omega)$ be a solution of $\mathcal{L}_\phi u = f$ in Ω , where $\phi \in C(\overline{\Omega})$ is a convex function satisfying $\phi = 0$ on $\partial \Omega$. There exists $\epsilon > 0$ depending only on ϵ_0, α_0 and n such that*

if $1 - \epsilon \leq \det D^2\phi \leq 1 + \epsilon$ in Ω , then for any $S_\phi(x_0, t_0) \subset \Omega_{\alpha_0}$ with $t_0 \leq \eta(\alpha_0)$ and $S_\phi(x_0, t_0) \cap G_\gamma(u, \Omega) \neq \emptyset$ we have

$$|G_{N\gamma}(u, \Omega) \cap S_\phi(x_0, t_0)| \geq \left\{ 1 - \epsilon_0 - C(N\gamma)^{-\tau} \left(\int_{S_\phi(x_0, \frac{t_0}{\alpha_0})} |f|^n dx \right)^{\frac{\tau}{n}} \right\} |S_\phi(x_0, t_0)|$$

for all $N \geq N_0$. Here $\eta(\alpha_0)$, C , τ and N_0 are constants depending only on α_0 and n .

Proof. Let $\theta > 1$ be the engulfing constant corresponding to $1/2 \leq \det D^2\phi \leq 3/2$ in Ω and so θ depends only on the dimension n . By Lemma 2.1, there exists $\eta(\alpha_0) = \eta(\alpha_0, n) > 0$ such that $S_\phi(x, \frac{\theta t}{\alpha_0}) \subset \Omega_{\frac{\alpha_0+1}{2}}$ for all $x \in \Omega_{\alpha_0}$ and $t \leq \eta(\alpha_0)$. We note that $\phi \in C^1(\Omega) \cap W^{2,n}(\Omega_{\frac{\alpha_0+1}{2}})$ as explained in the proof of Lemma 4.3.

Let T normalize $S_\phi(x_0, \frac{t_0}{\alpha_0})$ and $U := T(S_\phi(x_0, \frac{t_0}{\alpha_0}))$. For each $y \in T(\Omega)$, set

$$\tilde{\phi}(y) = |\det T|^{\frac{2}{n}} \left[\phi(T^{-1}y) - \phi(x_0) - \langle \nabla\phi(x_0), T^{-1}y - x_0 \rangle - \frac{t_0}{\alpha_0} \right] \text{ and } \tilde{u}(y) = \frac{1}{2\theta t_0} u(T^{-1}y).$$

We have U is normalized, $1 - \epsilon \leq \det D^2\tilde{\phi} \leq 1 + \epsilon$ in U and $\tilde{\phi} = 0$ on ∂U . Let $\bar{x} \in S_\phi(x_0, t_0) \cap G_\gamma(u, \Omega)$ and $\bar{y} = T\bar{x}$. Then

$$-\gamma d(x, \bar{x})^2 \leq u(x) - u(\bar{x}) - \nabla u(\bar{x}) \cdot (x - \bar{x}) \leq \gamma d(x, \bar{x})^2, \quad \forall x \in \Omega.$$

Hence by changing variables we get

(4.50)

$$-\gamma \frac{d(T^{-1}y, T^{-1}\bar{y})^2}{2\theta t_0} \leq \tilde{u}(y) - \tilde{u}(\bar{y}) - \nabla\tilde{u}(\bar{y}) \cdot (y - \bar{y}) \leq \gamma \frac{d(T^{-1}y, T^{-1}\bar{y})^2}{2\theta t_0}, \quad \forall y \in T(\Omega).$$

Since $\bar{x} \in S_\phi(x_0, t_0) \subset S_\phi(x_0, t_0/\alpha_0)$, we have $S_\phi(x_0, t_0/\alpha_0) \subset S_\phi(\bar{x}, \theta t_0/\alpha_0)$ by the engulfing property. It follows that $d(x, \bar{x})^2 \leq \theta t_0/\alpha_0$ for $x \in S_\phi(x_0, t_0/\alpha_0)$ yielding $d(T^{-1}y, T^{-1}\bar{y})^2 \leq \theta t_0/\alpha_0$ for all $y \in U$. Consequently,

$$-\frac{\gamma}{2\alpha_0} \leq \tilde{u}(y) - \tilde{u}(\bar{y}) - \nabla\tilde{u}(\bar{y}) \cdot (y - \bar{y}) \leq \frac{\gamma}{2\alpha_0} \quad \forall y \in U.$$

Let $v(y) := \frac{2\alpha_0}{\gamma} [\tilde{u}(y) - \tilde{u}(\bar{y}) - \nabla\tilde{u}(\bar{y}) \cdot (y - \bar{y})]$, for $y \in T(\Omega)$. Then $|v| \leq 1$ in U and by (4.50) we also have

$$|v(y)| \leq \frac{\alpha_0}{\theta t_0} d(T^{-1}y, T^{-1}\bar{y})^2 \leq \frac{C_n \alpha_0}{\theta} \tilde{d}(y, \bar{y})^2 \quad \forall y \in T(\Omega),$$

where $\tilde{d}(y, \bar{y})^2 := \tilde{\phi}(y) - \tilde{\phi}(\bar{y}) - \langle \nabla\tilde{\phi}(\bar{y}), y - \bar{y} \rangle = |\det T|^{2/n} d(T^{-1}y, T^{-1}\bar{y})^2$. Moreover

$$\text{trace}(\tilde{\Phi} D^2 v) = \frac{\alpha_0 |\det T|^{\frac{-2}{n}}}{\theta \gamma t_0} \text{trace}(\Phi(T^{-1}y) D^2 u(T^{-1}y)) = \frac{\alpha_0}{\theta \gamma t_0 |\det T|^{\frac{2}{n}}} f(T^{-1}y) =: \tilde{f}(y).$$

Notice that $\bar{y} \in U_{\alpha_0} = T(S_\phi(x_0, t_0))$ because $\bar{x} \in S_\phi(x_0, t_0)$. Thus we obtain from Lemma 4.2 with $\alpha := \alpha_0$ that

$$|G_N(v, T(\Omega)) \cap U_{\alpha_0}| \geq \left\{ 1 - C(N^{-\tau} \delta_0^\tau + \epsilon) \right\} |U_{\alpha_0}|$$

for any $N \geq N_0 = N_0(\alpha_0, n)$ and provided that $\|\tilde{\Phi} - \tilde{W}\|_{L^n(U_{\frac{\alpha_0+1}{2}})} \leq ((1 - \alpha_0)/4)^{\frac{2n}{1+(n-1)\gamma}}$, where δ_0 and \tilde{W} are as in (4.46). This together with Lemma 4.4 implies that there exists $\epsilon > 0$ depending only on ϵ_0, α_0 and n such that

$$\begin{aligned} |G_N(v, T(\Omega)) \cap U_{\alpha_0}| &\geq \left\{ 1 - \epsilon_0 - CN^{-\tau} \left(\int_U |\tilde{f}|^n dy \right)^{\frac{\tau}{n}} \right\} |U_{\alpha_0}| \\ &= \left\{ 1 - \epsilon_0 - C \left(\frac{\alpha_0}{\theta\gamma N} \right)^\tau (t_0 |\det T|^{\frac{2}{n}})^{-\tau} \left(\int_{S_\phi(x_0, \frac{t_0}{\alpha_0})} |f|^n dx \right)^{\frac{\tau}{n}} \right\} |U_{\alpha_0}| \\ &\geq \left\{ 1 - \epsilon_0 - C \left(\frac{\alpha_0}{\theta\gamma N} \right)^\tau \left(\int_{S_\phi(x_0, \frac{t_0}{\alpha_0})} |f|^n dx \right)^{\frac{\tau}{n}} \right\} |U_{\alpha_0}|. \end{aligned}$$

But since $U_{\alpha_0} = T(S_\phi(x_0, t_0))$ and $v(y) = \frac{\alpha_0}{\theta\gamma t_0} [u(T^{-1}y) - u(\bar{x}) - \langle \nabla u(\bar{x}), T^{-1}y - \bar{x} \rangle]$, the same calculations leading to (4.42) yield

$$G_N(v, T(\Omega)) \cap U_{\alpha_0} = T \left(G_{\frac{N\theta\gamma t_0 |\det T|^{\frac{2}{n}}}{\alpha_0}}(u, \Omega) \cap S_\phi(x_0, t_0) \right) \approx T \left(G_{\frac{N\theta\gamma}{\alpha_0}}(u, \Omega) \cap S_\phi(x_0, t_0) \right).$$

Therefore we obtain

$$\left| T \left(G_{\frac{N\theta\gamma}{\alpha_0}}(u, \Omega) \cap S_\phi(x_0, t_0) \right) \right| \geq \left\{ 1 - \epsilon_0 - C \left(\frac{\alpha_0}{\theta\gamma N} \right)^\tau \left(\int_{S_\phi(x_0, \frac{t_0}{\alpha_0})} |f|^n dx \right)^{\frac{\tau}{n}} \right\} |T(S_\phi(x_0, t_0))|.$$

By setting $N' = N\theta/\alpha_0$, we can rewrite this as

$$|G_{N'\gamma}(u, \Omega) \cap S_\phi(x_0, t_0)| \geq \left\{ 1 - \epsilon_0 - C \left(\frac{1}{\gamma N'} \right)^\tau \left(\int_{S_\phi(x_0, \frac{t_0}{\alpha_0})} |f|^n dx \right)^{\frac{\tau}{n}} \right\} |S_\phi(x_0, t_0)|$$

for any $N' \geq N_0 = N_0(\alpha_0, n)$. □

4.2. $W^{2,p}$ estimate. In this subsection we will use the density estimates established in Subsection 4.1 to derive interior $W^{2,p}$ -estimates for solution u of the linearized equation $\mathcal{L}_\phi u = f$ when $f \in L^q(\Omega)$ for some $q > n$. We begin with the following key result which gives a solution to the conjecture in [GT].

Theorem 4.6. *Let Ω be a normalized convex domain and $u \in W_{loc}^{2,n}(\Omega)$ be a solution of $\mathcal{L}_\phi u = f$ in Ω , where $\phi \in C(\overline{\Omega})$ is a convex function satisfying $\phi = 0$ on $\partial\Omega$. Let $p > 1$, $\max\{n, p\} < q < \infty$ and let $0 < \alpha < 1$. Then there exist positive constants ϵ and C depending only on p, q, α and n such that if $1 - \epsilon \leq \det D^2\phi \leq 1 + \epsilon$, we have*

$$\|D^2u\|_{L^p(\Omega_\alpha)} \leq C(\|u\|_{L^\infty(\Omega)} + \|f\|_{L^q(\Omega)}).$$

Proof. We first observe that by working with the function $v := \frac{\epsilon u}{\epsilon \|u\|_{L^\infty(\Omega)} + \|f\|_{L^q(\Omega)}}$ instead of u , it is enough to show that there exist $\epsilon, C > 0$ depending only on p, q, α and n such that if $1 - \epsilon \leq \det D^2\phi \leq 1 + \epsilon$, $\|u\|_{L^\infty(\Omega)} \leq 1$ and $\|f\|_{L^q(\Omega)} \leq \epsilon$, then

$$(4.51) \quad \|D^2u\|_{L^p(\Omega_\alpha)} \leq C.$$

Note also that $u \in C^1(\Omega)$ as a consequence of $C_{loc}^{1,\alpha}$ estimates in [GN, Theorem 4.5].

Let $\alpha_0 := \frac{\alpha+1}{2}$ and $N_0 = N_0(\alpha_0, n)$ be the largest of the constants in Lemma 4.3 and Lemma 4.5. Fix $M \geq N_0$ so that $\frac{1}{C_0^{1/\gamma}(M^{1/\gamma}-1)} \leq \frac{1-\alpha_0}{2}$ and $(cM^{\frac{q-p}{2p}})^{\frac{1}{n-1}} \geq \frac{\text{diam}(\Omega)}{\sqrt{\eta_0}}$, where γ, C_0 are given by Lemma 2.1 and c is given by Lemma 2.7 when $\lambda = 1/2$ and $\Lambda = 3/2$. Next select $0 < \epsilon_0 < 1/2$ such that

$$M^q \sqrt{2\epsilon_0} = \frac{1}{2}$$

and $\epsilon = \epsilon(\epsilon_0, \alpha_0, n) = \epsilon(p, q, \alpha, n)$ be the smallest of the constants in Lemma 4.3 and Lemma 4.5. With this choice of ϵ , we are going to show that (4.51) holds. Applying Lemma 4.3 to the function u and using $\|f\|_{L^q(\Omega)} \leq \epsilon$ we obtain

$$|S_\phi(x_0, t_0) \cap G_{\frac{M}{t_0}}(u, \Omega)| \geq (1 - \epsilon_0 - C\epsilon^\tau) |S_\phi(x_0, t_0)|$$

as long as $S_\phi(x_0, \frac{t_0}{\alpha_0}) \subset \Omega_{\frac{\alpha_0+1}{2}}$, where $C = C(p, \alpha, n)$ and $\tau = \tau(\alpha, n)$. By taking ϵ even smaller if necessary we can assume $C\epsilon^\tau < \epsilon_0$. Then it follows from the above inequality that

$$(4.52) \quad |S_\phi(x_0, t_0) \setminus G_{\frac{M}{t_0}}(u, \Omega)| \leq 2\epsilon_0 |S_\phi(x_0, t_0)| \quad \text{for any } S_\phi(x_0, \frac{t_0}{\alpha_0}) \subset \Omega_{\frac{\alpha_0+1}{2}}.$$

Let $\eta(\alpha_0) > 0$ be given by Lemma 4.5 ensuring in particular that $S_\phi(x, \frac{t}{\alpha_0}) \subset \Omega_{\frac{\alpha_0+1}{2}}$ for all $x \in \Omega_{\alpha_0}$ and $t \leq \eta(\alpha_0)$. Assume $\alpha_2 < \alpha_1 < \alpha_0$ are such that there exist $\eta_2 < \eta_1 \leq \eta(\alpha_0)$ with the property: if $x \in \Omega_{\alpha_2}$ and $t \leq \eta_2$ then $S_\phi(x, t) \subset \Omega_{\alpha_1}$; and if $x \in \Omega_{\alpha_1}$ and $t \leq \eta_1$ then $S_\phi(x, t) \subset \Omega_{\alpha_0}$. With these choices and for $1/h \leq \eta_2$, by using (4.52) and the same arguments leading to (3.25) we obtain: for any $x_0 \in \Omega_{\alpha_2} \setminus G_{hM}(u, \Omega)$ there is $t_{x_0} \leq 1/h$ such that

$$(4.53) \quad |(\Omega_{\alpha_2} \setminus G_{hM}(u, \Omega)) \cap S_\phi(x_0, t_{x_0})| = 2\epsilon_0 |S_\phi(x_0, t_{x_0})|.$$

We now claim that (4.53) implies

$$(4.54) \quad S_\phi(x_0, t_{x_0}) \subset (\Omega_{\alpha_1} \setminus G_h(u, \Omega)) \cup \{x \in \Omega_{\alpha_0} : \mathcal{M}_\mu((f/\det D^2\phi)^n)(x) > (c^*Mh)^n\},$$

where $c^* := (\frac{\epsilon_0}{C})^{1/\tau}$ and $\mu := M\phi$. Otherwise, and since $x_0 \in \Omega_{\alpha_2}$ and $t_{x_0} \leq 1/h \leq \eta_2$, we have that $S_\phi(x_0, t_{x_0}) \subset \Omega_{\alpha_1}$ and there exists $\bar{x} \in S_\phi(x_0, t_{x_0}) \cap G_h(u, \Omega)$ such that $\mathcal{M}_\mu((f/\det D^2\phi)^n)(\bar{x}) \leq (c^*Mh)^n$. Note also that $t_{x_0} \leq \eta(\alpha_0)$ and due to our assumption on ϕ the measure μ is comparable to the Lebesgue measure. Then by Lemma 4.5 applied to u we get

$$|S_\phi(x_0, t_{x_0}) \cap G_{hM}(u, \Omega)| > (1 - 2\epsilon_0) |S_\phi(x_0, t_{x_0})|$$

yielding

$$\left| \left(\Omega_{\alpha_2} \setminus G_{hM}(u, \Omega) \right) \cap S_\phi(x_0, t_{x_0}) \right| \leq \left| S_\phi(x_0, t_{x_0}) \setminus G_{hM}(u, \Omega) \right| < 2\epsilon_0 |S_\phi(x_0, t_{x_0})|.$$

This is a contradiction with (4.53) and so (4.54) is proved. We infer from (4.53), (4.54) and [G, Theorem 6.3.3] that

$$(4.55) \quad \begin{aligned} & |\Omega_{\alpha_2} \setminus G_{hM}(u, \Omega)| \\ & \leq \sqrt{2\epsilon_0} \left[|\Omega_{\alpha_1} \setminus G_h(u, \Omega)| + \left| \{x \in \Omega_{\alpha_0} : \mathcal{M}_\mu((f/\det D^2\phi)^n)(x) > (c^*Mh)^n\} \right| \right], \end{aligned}$$

as long as $\alpha_2 < \alpha_1 < \alpha_0$ are such that $\eta_2 < \eta_1 \leq \eta(\alpha_0)$, and $1/h \leq \eta_2$.

For $k = 0, 1, \dots$, set

$$a_k := |\Omega_{\alpha_k} \setminus G_{M^k}(u, \Omega)| \quad \text{and} \quad b_k := \left| \{x \in \Omega_{\alpha_0} : \mathcal{M}_\mu((f/\det D^2\phi)^n)(x) > (c^*MM^k)^n\} \right|,$$

where α_k will be defined inductively in the sequel. First fix α_1 so that

$$\frac{3\alpha_0 - 1}{2} < \alpha_1 < \alpha_0 \quad \text{and} \quad \eta_1 := C_0(\alpha_0 - \alpha_1)^\gamma \leq \eta(\alpha_0).$$

By taking M even larger if necessary, we can assume that $1/M < \eta_1$. Let $h = M$, and set $\alpha_2 = \alpha_1 - \frac{1}{(C_0M)^{1/\gamma}}$. Then $\frac{1}{h} = \frac{1}{M} = C_0(\alpha_1 - \alpha_2)^\gamma =: \eta_2$, and so from Lemma 2.1 and (4.55) we get $a_2 \leq \sqrt{2\epsilon_0}(a_1 + b_1)$. Next let $h = M^2$ and $\alpha_3 = \alpha_2 - \frac{1}{(C_0M^2)^{1/\gamma}}$, so $\frac{1}{h} = C_0(\alpha_2 - \alpha_3)^\gamma =: \eta_3$. Then $a_3 \leq \sqrt{2\epsilon_0}(a_2 + b_2) \leq 2\epsilon_0 a_1 + 2\epsilon_0 b_1 + \sqrt{2\epsilon_0} b_2$. Continuing in this way we conclude that

$$a_{k+1} \leq (\sqrt{2\epsilon_0})^k a_1 + \sum_{i=1}^k (\sqrt{2\epsilon_0})^{(k+1)-i} b_i \quad \text{for } k = 1, 2, \dots$$

On the other hand, $\alpha_{k+1} = \alpha_1 - \sum_{j=1}^k \frac{1}{(C_0M^j)^{1/\gamma}} \geq \frac{3\alpha_0 - 1}{2} - \frac{1}{C_0^{1/\gamma}(M^{1/\gamma-1})} \geq 2\alpha_0 - 1 = \alpha$ by our choice of α_1, M and α_0 . Therefore for every $k \geq 1$,

$$(4.56) \quad |\Omega_\alpha \setminus G_{M^{k+1}}(u, \Omega)| \leq |\Omega_{\alpha_{k+1}} \setminus G_{M^{k+1}}(u, \Omega)| \leq (\sqrt{2\epsilon_0})^k a_1 + \sum_{i=1}^k (\sqrt{2\epsilon_0})^{(k+1)-i} b_i.$$

Next let $\Theta(u)$ be the function defined by (2.8). We claim that $\Theta(u) \in L^p(\Omega_\alpha)$ and

$$(4.57) \quad \|\Theta(u)\|_{L^p(\Omega_\alpha)} \leq C(p, q, \alpha, n).$$

Indeed since $u \in C^1(\Omega)$, it is easy to see that $\Theta(u)$ is lower semicontinuous in Ω_α and so measurable there. Moreover, we have

$$\begin{aligned}
 \int_{\Omega_\alpha} |\Theta(u)|^p dx &= p \int_0^\infty t^{p-1} |\{x \in \Omega_\alpha : \Theta(u)(x) > t\}| dt \\
 &= p \int_0^{M^{\frac{q}{p}}} t^{p-1} |\{x \in \Omega_\alpha : \Theta(u)(x) > t\}| dt + p \sum_{k=1}^\infty \int_{M^{\frac{qk}{p}}}^{M^{\frac{q(k+1)}{p}}} t^{p-1} |\{x \in \Omega_\alpha : \Theta(u)(x) > t\}| dt \\
 &\leq |\Omega_\alpha| M^q + (M^q - 1) \sum_{k=1}^\infty M^{qk} |\{x \in \Omega_\alpha : \Theta(u)(x) > M^{\frac{qk}{p}}\}| \\
 &\leq |\Omega_\alpha| M^q + (M^q - 1) \left[\sum_{k=1}^\infty M^{qk} |\Omega_\alpha \setminus D_\alpha^{\alpha, \frac{k(q-p)}{(cM^{\frac{k(q-p)}})^{\frac{1}{n-1}}}}| + \sum_{k=1}^\infty M^{qk} |\Omega_\alpha \setminus G_{M^k}(u, \Omega)| \right] \\
 &\leq |\Omega_\alpha| M^q + (M^q - 1) \left[\frac{|\Omega|}{(C_n \epsilon)^2} c^{\ln \sqrt{C_n \epsilon}} \sum_{k=1}^\infty M^{k(q + (\frac{q}{p}-1)\frac{\ln \sqrt{C_n \epsilon}}{c})} + \sum_{k=1}^\infty M^{qk} |\Omega_\alpha \setminus G_{M^k}(u, \Omega)| \right],
 \end{aligned}$$

where we used (2.9) with $\kappa = q/p > 1$ and $\beta = M^k$ in the second inequality and used (2.10) in the last inequality. Since $\epsilon > 0$ is small, the first summation in the last expression is finite and hence (4.57) will follow if we can show that $\sum_{k=1}^\infty M^{kq} |\Omega_\alpha \setminus G_{M^k}(u, \Omega)| \leq C$. For this, let us employ (4.56) to obtain

$$\begin{aligned}
 \sum_{k=1}^\infty M^{kq} |\Omega_\alpha \setminus G_{M^k}(u, \Omega)| &\leq a_1 \sum_{k=1}^\infty M^{kq} (\sqrt{2\epsilon_0})^{k-1} + \sum_{k=1}^\infty \sum_{i=0}^{k-1} M^{kq} (\sqrt{2\epsilon_0})^{k-i} b_i \\
 &= \frac{a_1}{\sqrt{2\epsilon_0}} \sum_{k=1}^\infty (M^q \sqrt{2\epsilon_0})^k + \sum_{i=0}^\infty \sum_{k=i+1}^\infty M^{(k-i)q} (\sqrt{2\epsilon_0})^{k-i} M^{iq} b_i \\
 &= \frac{a_1}{\sqrt{2\epsilon_0}} \sum_{k=1}^\infty (M^q \sqrt{2\epsilon_0})^k + \left[\sum_{j=1}^\infty (M^q \sqrt{2\epsilon_0})^j \right] \left[\sum_{i=0}^\infty M^{iq} b_i \right] \\
 &= \frac{a_1}{\sqrt{2\epsilon_0}} \sum_{k=1}^\infty 2^{-k} + \left[\sum_{j=1}^\infty 2^{-j} \right] \left[\sum_{i=0}^\infty M^{iq} b_i \right] = \frac{a_1}{\sqrt{2\epsilon_0}} + \sum_{i=0}^\infty M^{iq} b_i.
 \end{aligned}$$

But as $f^n \in L^{\frac{q}{n}}(\Omega)$ and $q > n$, we have from Theorem 2.2 that

$$\int_{\Omega_{\alpha_0}} \left| \mathcal{M}_\mu \left(\left(\frac{f}{\det D^2 \phi} \right)^n \right) (x) \right|^{\frac{q}{n}} d\mu(x) \leq C_n \int_\Omega \left| \frac{f}{\det D^2 \phi} \right|^q d\mu(y) \leq C$$

implying $\sum_{i=0}^\infty (M^n)^{i\frac{q}{n}} b_i \leq C$. Thus $\sum_{k=1}^\infty M^{kq} |\Omega_\alpha \setminus G_{M^k}(u, \Omega)| \leq C$ and claim (4.57) is proved.

It follows from (4.57) and [CC, Proposition 1.1] that $D^2 u \in L^p(\Omega_\alpha)$ and $\|D^2 u\|_{L^p(\Omega_\alpha)} \leq 4\|\Theta(u)\|_{L^p(\Omega_\alpha)} \leq C(p, q, \alpha, n)$. This gives (4.51) as desired and the proof is complete. \square

We are finally in a position to prove the main result of the paper, Theorem 1.1.

Proof of Theorem 1.1. Let $\epsilon = \epsilon(p, q, n)$ be the constant given by Theorem 4.6 corresponding to $\alpha = 1/2$. Let $x \in \Omega'$ and suppose a section $S = S_\phi(x, \delta) \Subset \Omega$ is such that $|g(z) - g(x)| \leq \lambda\epsilon$, for each $z \in S$. Then by the property of sections [G, Theorem 3.3.8], we have

$$(4.58) \quad B(x, K_1\delta) \subset S \subset B(x, K_2\delta^b),$$

with K_1, K_2, b positive constants depending only on λ, Λ and n . Let $Tx = Ax + b$ be an affine map normalizing S and consider the following functions on $\tilde{\Omega} := T(S)$:

$$\begin{aligned} \tilde{\phi}(y) &:= \frac{|\det A|^{\frac{2}{n}}}{g(x)^{\frac{1}{n}}} \left[\phi(T^{-1}y) - \phi(x) - \nabla\phi(x) \cdot (T^{-1}y - x) - \delta \right], \\ \text{and } \tilde{u}(y) &:= |\det A|^{\frac{2}{n}} g(x)^{\frac{n-1}{n}} u(T^{-1}y). \end{aligned}$$

We have $\tilde{\Omega}$ is normalized, $\tilde{\phi} = 0$ on $\partial\tilde{\Omega}$ and $D^2\tilde{\phi}(y) = \frac{|\det A|^{\frac{2}{n}}}{g(x)^{\frac{1}{n}}} (A^{-1})^t D^2\phi(T^{-1}y) A^{-1}$.

Thus $\det D^2\tilde{\phi}(y) = \frac{g(T^{-1}y)}{g(x)} =: \tilde{g}(y)$ and if $\tilde{\Phi}(y)$ is the cofactor matrix of $D^2\tilde{\phi}(y)$, then

$$\mathcal{L}_{\tilde{\phi}}\tilde{u}(y) = \text{trace}(\tilde{\Phi}(y) D^2\tilde{u}(y)) = \text{trace}(\Phi(T^{-1}y) D^2u(T^{-1}y)) = f(T^{-1}y) =: \tilde{f}(y) \quad \text{in } \tilde{\Omega}.$$

Moreover since $g(x) - \lambda\epsilon \leq g(z) \leq g(x) + \lambda\epsilon$ for $z \in S$ and $g \geq \lambda$, we get

$$1 - \epsilon \leq 1 - \frac{\epsilon\lambda}{g(x)} \leq \tilde{g}(y) \leq 1 + \frac{\lambda\epsilon}{g(x)} \leq 1 + \epsilon \quad \text{for } y \in \tilde{\Omega}.$$

Therefore, we can apply Theorem 4.6 to obtain

$$(4.59) \quad \left(\int_{\tilde{\Omega}_{1/2}} |D^2\tilde{u}(y)|^p dy \right)^{1/p} \leq C(p, q, n) (\|\tilde{u}\|_{L^\infty(\tilde{\Omega})} + \|\tilde{f}\|_{L^q(\tilde{\Omega})}) \\ = C(|\det A|^{\frac{2}{n}} g(x)^{\frac{n-1}{n}} \|u\|_{L^\infty(S)} + |\det A|^{\frac{1}{q}} \|f\|_{L^q(S)}).$$

By the definition of \tilde{u} we have $D^2u(z) = |\det A|^{\frac{2}{n}} g(x)^{\frac{1-n}{n}} A^t D^2\tilde{u}(Tz) A$ in S , and consequently $\|D^2u\|_{L^p(S_{1/2})} \leq \|A\|^2 |\det A|^{-\frac{2}{n} + \frac{1}{p}} g(x)^{\frac{1-n}{n}} \|D^2\tilde{u}\|_{L^p(\tilde{\Omega}_{1/2})}$ where $S_{1/2} := S_\phi(x, \delta/2)$.

Notice that $|\det A| \approx \delta^{-n/2}$ by the normalization, and $\|A\| \leq C\delta^{-1}$ by the fact $AB(x, K_1\delta) + b \subset B_n(0)$ following from (4.58). Hence we deduce from (4.59) that

$$(4.60) \quad \left(\int_{S_{1/2}} |D^2u(z)|^p dz \right)^{1/p} \leq C\|A\|^2 |\det A|^{\frac{-1}{p}} (\|u\|_{L^\infty(S)} + |\det A|^{\frac{1}{q} - \frac{2}{n}} g(x)^{\frac{1-n}{n}} \|f\|_{L^q(S)}) \\ \leq C\delta^{\frac{n}{2p} - 2} \|u\|_{L^\infty(\Omega)} + C\delta^{\frac{n}{2p} - \frac{n}{2q} - 1} \|f\|_{L^q(\Omega)},$$

where C depends only on p, q, λ, Λ and n .

Now since $\Omega' \Subset \Omega$, we can pick δ small depending only on the parameters $\lambda, \Lambda, n, \text{dist}(\Omega', \partial\Omega)$ and the modulus of continuity of g such that for each $x \in \Omega'$ we have $B(x, K_2\delta^b) \Subset \Omega$ and $|g(z) - g(x)| \leq \lambda\epsilon$ in $B(x, K_2\delta^b)$. Next select a finite

covering of Ω' by balls $\{B(x_j, K_1\delta)\}_{j=1}^N$ with $x_j \in \Omega'$, then the desired inequality follows by adding (4.60) over $\{S_\phi(x_j, \delta/2)\}_{j=1}^N$. \square

In this paper we have chosen to work with strong solutions in $W_{loc}^{2,n}(\Omega)$ in order to reveal direct calculations. However, the interior $W^{2,p}$ estimates in Theorem 4.6 and Theorem 1.1 can be derived for viscosity solutions of $\mathcal{L}_\phi u = f$ by modifying slightly the definition of the set $G_M(u, \Omega)$ and following our arguments. For this purpose we note that the Caffarelli-Gutiérrez interior Hölder estimates, which were used in Lemma 4.1, still hold for viscosity solutions as observed by Trudinger and Wang in [TW4].

REFERENCES

- [C1] L. A. Caffarelli. *Interior a priori estimates for solutions of fully nonlinear equations*. Ann. of Math. **130**, no. 1, 189–213, 1989.
- [C2] L. A. Caffarelli. *A localization property of viscosity solutions to the Monge–Ampère equation and their strict convexity*. Ann. of Math. **131**, 129–134, 1990.
- [C3] L. A. Caffarelli. *Interior $W^{2,p}$ estimates for solutions to the Monge–Ampère equation*. Ann. of Math. **131**, 135–150, 1990.
- [C4] L. A. Caffarelli. *Some regularity properties of solutions of Monge–Ampère equation*. Comm. Pure Appl. Math. **44**, 965–969, 1991.
- [CC] L. A. Caffarelli and X. Cabré. *Fully nonlinear elliptic equations*. American Mathematical Society Colloquium Publications, volume 43, 1995.
- [CG1] L. A. Caffarelli and C. E. Gutiérrez. *Real analysis related to the Monge–Ampère equation*. Trans. Amer. Math. Soc. **348**, no. 3, 1075–1092, 1996.
- [CG2] L. A. Caffarelli and C. E. Gutiérrez. *Properties of the solutions of the linearized Monge–Ampère equation*. Amer. J. Math. **119**, no. 2, 423–465, 1997.
- [CNP] M. J. Cullen, J. Norbury and R. J. Purser. *Generalized Lagrangian solutions for atmospheric and oceanic flows*. SIAM J. Appl. Math. **51**, no. 1, 20–31, 1991.
- [dPF] G. De Phillips and A. Figalli. *$W^{2,1}$ regularity for solutions of the Monge–Ampère equation*. Preprint, 2011.
- [D1] S. K. Donaldson. *Scalar curvature and stability of toric varieties*. J. Differential Geom. **62**, no. 2, 289–349, 2002.
- [D2] S. K. Donaldson. *Interior estimates for solutions of Abreu’s equation*. Collect. Math. **56**, no. 2, 103–142, 2005.
- [D3] S. K. Donaldson. *Extremal metrics on toric surfaces: a continuity method*. J. Differential Geom. **79**, no. 3, 389–432, 2008.
- [D4] S. K. Donaldson. *Constant scalar curvature metrics on toric surfaces*. Geom. Funct. Anal. **19**, no. 1, 83–136, 2009.
- [Es] L. Escauriaza. *$W^{2,n}$ a priori estimates for solutions to fully non-linear equations*. Indiana University Mathematics Journal **42**, no. 2, 413–423, 1993.
- [E] L. C. Evans. *Some estimates for non divergence structure, second order elliptic equations*. Trans. Amer. Math. Soc. **287**, no. 2, 701–712, 1985.
- [GiT] D. Gilbarg and N.S. Trudinger. *Elliptic partial differential equations of second order*. Springer–Verlag, New York, 2001.
- [G] C. E. Gutiérrez. *The Monge–Ampère equation*. Birkhäuser, Boston, 2001.
- [GH] C. E. Gutiérrez and Q. Huang. *Geometric properties of the sections of solutions to the Monge–Ampère equation*. Trans. Amer. Math. Soc. **352**, 4381–4396, 2000.

- [GN] C. E. Gutiérrez and T.V. Nguyen. *Interior gradient estimates for solutions to the linearized Monge-Ampère equation*. Adv. Math., **228**, 2034–2070, 2011.
- [GT] C. E. Gutiérrez and F. Tournier. *$W^{2,p}$ -estimates for the linearized Monge-Ampère equation*. Trans. Amer. Math. Soc. **358**, no. 11, 4843–4872, 2006.
- [H] Q. Huang. *Sharp regularity results on second derivatives of solutions to the Monge-Ampère equation with VMO type data*. Comm. Pure Appl. Math. **62**, no. 5, 677–705, 2009.
- [LS1] N. Q. Le and O. Savin. *Boundary regularity for solutions to the linearized Monge-Ampère equations*. Preprint, 2011.
- [LS2] N. Q. Le and O. Savin. *Some minimization problems in the class of convex functions with prescribed determinant*. Preprint, 2011.
- [L] F.-H. Lin. *Second derivative L^p -estimates for elliptic equations of nondivergent type*. Proc. Amer. Math. Soc. **96**, no. 3, 447–451, 1986.
- [Lo] G. Loeper. *A fully nonlinear version of the incompressible Euler equations: the semigeostrophic system*. SIAM J. Math. Anal. **38**, no. 3, 795–823, 2006.
- [PT] C. Pucci and G. Talenti. *Elliptic (second-order) partial differential equations with measurable coefficients and approximating integral equations*. Adv. Math., **19**, no. 1, 48–105, 1976.
- [S1] O. Savin. *A Liouville theorem for solutions to the linearized Monge-Ampère equation*. Discrete Contin. Dyn. Syst. **28**, no. 3, 865–873, 2010.
- [S2] O. Savin. *Global $W^{2,p}$ estimates for the Monge-Ampère equations*. Preprint, 2010.
- [Sc] R. Schneider. *Convex bodies: The Brunn-Minkowski theory*. Cambridge University Press, Cambridge, 1993.
- [St] E.M. Stein. *Singular integrals and differentiability properties of functions*. Princeton University Press, Princeton, N.J., 1970.
- [Sw] A. Świech. *$W^{1,p}$ -interior estimates for solutions of fully nonlinear, uniformly elliptic equations*. Adv. Differential Equations **2**, no. 6, 1005–1027, 1997.
- [TiW] G.-J. Tian and X.-J. Wang. *A class of Sobolev type inequalities*. Methods and Applications of Analysis **15**, no. 2, 263–276, 2008.
- [T] N. Trudinger. *Glimpses of nonlinear partial differential equations in the twentieth century: A priori estimates and the Bernstein problem*. Challenges for the 21st century (Singapore, 2000), 196–212, World Sci. Publ., River Edge, NJ, 2001.
- [TW1] N. Trudinger and X.-J. Wang. *The Bernstein problem for affine maximal hypersurfaces*. Invent. Math. **140**, no. 2, 399–422, 2000.
- [TW2] N. Trudinger and X.-J. Wang. *The affine plateau problem*. J. Amer. Math. Soc. **18**, 253–289, 2005.
- [TW3] N. Trudinger and X.-J. Wang. *Boundary regularity for Monge-Ampère and affine maximal surface equations*. Ann. of Math. **167**, 993–1028, 2008.
- [TW4] N. Trudinger and X.-J. Wang. *The Monge-Ampère equation and its geometric applications*. Handbook of geometric analysis. No. 1, Adv. Lect. Math., 7, Int. Press, Somerville, MA, 467–524, 2008.
- [U] N. Uralt'ceva. *The impossibility of $W^{2,q}$ estimates for multidimensional elliptic equations with discontinuous coefficients*. (Russian) Zap. Naun. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **5**, 250–254, 1967.
- [WL] L. Wang. *A geometric approach to the Calderón-Zygmund estimates*. Acta Math. Sin. (Engl. Ser.) **19**, no. 2, 381–396, 2003.
- [W] X.-J. Wang. *Some counterexamples to the regularity of Monge-Ampère equations*. Proc. Amer. Math. Soc. **123**, no. 3, 841–845, 1995.
- [Z1] B. Zhou. *The first boundary value problem for Abreu's equation*. Preprint, 2010.
- [Z2] B. Zhou. *The Bernstein theorem for a class of fourth order equations*. Calc. Var. Partial Differential Equations, doi 10.1007/s00526-011-0401-3.

CRISTIAN GUTIÉRREZ , TEMPLE UNIVERSITY, DEPARTMENT OF MATHEMATICS, PHILADELPHIA, PA 19122, USA.

E-mail address: gutierre@temple.edu

TRUYEN NGUYEN, THE UNIVERSITY OF AKRON, DEPARTMENT OF MATHEMATICS, AKRON, OH 44325, USA.

E-mail address: tnguyen@uakron.edu