

Author Manuscript

Title: Concise Syntheses of Bis-Strychnos Alkaloids (–)-Sungucine, (–)-Isosungucine, and (–)-Strychnogucine B from (–)-Strychnine

Authors: Senzhi Zhao; Christiana Teijaro; Heng Chen; Gopal Sirasani; Shivaiah Vaddypally; Michael Zdilla; Graham Dobereiner; Rodrigo Andrade

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record.

To be cited as: 10.1002/chem.201602663

Link to VoR: <http://dx.doi.org/10.1002/chem.201602663>

Concise Syntheses of Bis-*Strychnos* Alkaloids (–)-Sungucine, (–)-Isosungucine, and (–)-Strychnogucine B from (–)-Strychnine


Senzhi Zhao, Christiana N. Teijaro, Heng Chen, Gopal Sirasani, Shivaiah Vaddypally, Michael J. Zdilla, Graham E. Dobereiner, and Rodrigo B. Andrade*

Abstract: We report the first chemical syntheses of complex, bis-*Strychnos* alkaloids (–)-sungucine (**1**), (–)-isosungucine (**2**), and (–)-strychnogucine B (**3**) from (–)-strychnine (**4**). Key steps included (1) the Polonovski–Potier activation of strychnine *N*-oxide; (2) a biomimetic Mannich coupling to forge the signature C23–C5' bond that joins two monoterpene indole monomers; and, (3) a sequential HBr/NaBH₃CN-mediated reduction to fashion the ethylidene moieties in **1–3**. DFT calculations were employed to rationalize the regiochemical course of reactions involving strychnine congeners.

The chemical synthesis of complex natural products remains a vibrant and dynamic enterprise within the rich field of organic chemistry. Many natural products are represented in our current pharmacopeia, including those in the bis-indole alkaloid class.¹ (–)-Sungucine (**1**) was isolated by Angenot and co-workers in 1979 from the roots of *Strychnos icaja*, a liana native only to central Africa (Scheme 1).² The structure of **1** was rigorously established by single-crystal X-ray analysis.³ Reinvestigation of *Strychnos icaja* by Fr d rich and co-workers in the 1990s resulted in the discovery of novel sungucine congeners (–)-isosungucine (**2**) and (–)-strychnogucine B (**3**).⁴

The structurally unique bis-*Strychnos* alkaloids **1–3** feature a common C23–C5' bond yet differ in oxidation states at C18 and C18' (Scheme 1). (–)-Sungucine (**1**) possesses anticancer activity in the low μ M range, as confirmed through NCI's *in vitro* 60 human tumor cell line screen.⁵ Pharmacological evaluation of **1** revealed it induced apoptosis in both leukemia and colon cancer cells via a p53-independent mechanism.⁶ In addition, **2** and **3** were found to possess potent and selective antiplasmodial activities against *Plasmodium falciparum* strains.^{4,7} Herein, we report the first enantiospecific chemical syntheses of bioactive, bis-*Strychnos* alkaloids (–)-sungucine (**1**), (–)-isosungucine (**2**), and (–)-strychnogucine B (**3**) from commercially available and inexpensive (–)-strychnine (**4**).

In our retrosynthetic analysis (Scheme 1), we envisioned (–)-sungucine (**1**) and (–)-isosungucine (**2**) could be synthesized from (–)-strychnogucine B (**3**) by base-mediated ring-opening and deoxygenation of the ethylidene moieties. To make (–)-**3** in a convergent manner, one can logically disconnect the C23–C5' bond, which could arise from the biomimetic Mannich coupling of a northern C23-synthon with a southern C5'-synthon. As both synthons are structurally homologous to strychnine (**4**), we reasoned the most step-efficient syntheses of **1–3** must start with inexpensive, commercially available (–)-strychnine (**4**).




Scheme 1. Structures of (–)-sungucine (**1**), (–)-isosungucine (**2**), (–)-strychnogucine B (**3**) and retrosynthetic analysis leading to northern and southern synthons derived from (–)-strychnine (**4**).

As outlined in Scheme 2, our syntheses of bis-*Strychnos* alkaloids **1–3** began with the construction of northern fragment **5** (i.e., TBS-protected isostrychnine). In order to readily access **5** on a multi-gram scale, we needed a reliable and effective method for converting strychnine into isostrychnine. While literature reports of this transformation are known,⁸ we were not satisfied with the results. Ultimately, we found that when (–)-strychnine (**4**) was first treated with 1,8-diazabicyclo[5.4.0]undecene (DBU) at 200 °C in *N*-methyl-2-pyrrolidone (NMP) then TBSCl and imidazole at rt, we routinely isolated **5** in 50% overall yield, in addition to minor regioisomer **6** (26% yield) on gram scale. Experimental results were consistent with DFT calculations inasmuch as only **5** and 13-*epi*-isostrychnine (**6**) were isolated out of the three possible products (see Table S1).

[a] S. Zhao, C. N. Teijaro, H. Chen, G. Sirasani, S. Vaddypally, M. J. Zdilla, G. E. Dobereiner, R. B. Andrade
Department of Chemistry
Temple University
1901 N. 13th St. Philadelphia, Pa 19122
E-mail: randrade@temple.edu

Supporting information for this article is given via a link at the end of the document.



Scheme 2. Syntheses of Northern fragment 5 and Southern fragment 9 from (-)-strychnine (4).

The next stage of the synthesis enlisted the venerable Polonovski–Potier reaction⁹ to regioselectively generate and trap C5–N4 iminium ion I (Figure 1) to access southern fragment 9 (Scheme 2). Thus, (-)-4 was oxidized with H₂O₂ to afford strychnine *N*-oxide in 98% yield. Reaction with trifluoroacetic anhydride (TFAA) then KCN gave an intractable mixture of regioisomers. Replacing KCN with NaHCO₃ (aq.) furnished pseudostrychnine (7) and regioisomeric carbinolamine 8 in a 2.2:1 ratio, respectively, and 71% yield. The structure of 7 was confirmed by single-crystal X-ray analysis. The use of more basic and nucleophilic KOH was able to invert the ratio of products and delivered the desired regioisomer 8 in 52% yield, along with 40% of 7. Spectral (i.e., ¹H NMR) analysis of major carbinolamine 8 revealed it was a mixture of epimers at C5. Treatment of this mixture with 10% MeOH/CH₂Cl₂ effected *N,O*-aminal ether equilibration to afford the requisite southern fragment 9 in quantitative yield. The structure of 9 was rigorously established by single-crystal X-ray analysis (Scheme 2).

To rationalize the regiochemical course of the Polonovski–Potier reaction, we employed DFT analysis (mPW1PW91/cc-pvdz) to determine relative free energies of three iminium cations that may form upon trifluoroacetate elimination (Figure 1). The carbinolamine derived from the lowest-energy iminium considered III was not observed in experiment; meanwhile, isolated regioisomer 7 was derived from the highest-energy iminium II. Besides the error inherent in the chosen DFT method, the effects of ion pairing¹⁰ may dictate the stability of I–III. Nonetheless, at this stage the experimental results cannot be understood solely by comparing the relative computed energies of I–III.




Figure 1. Relative DFT energies for iminium ions I–III.

Further DFT analysis afforded the minimized structure of parent *N*-acyloxyammonium ion IV (Table S2). IV lacks a hydrogen *anti*-periplanar to the N–O_{acyl} bond (Figure S1A–C), a strict prerequisite for the E2 pathway expected in the

Polonovski–Potier mechanism. The stereochemistry between C5–H_b and N4–O_{acyl} bonds (i.e., H_b–C5–N4–O dihedral=148.3°, Figures S1A and 2) are closest to being *anti*-periplanar. The formation of 8 may follow an E2-like mechanism via elimination from IV to I.

A natural population analysis (NPA)¹¹ of IV (Figure 2) indicates that C7 is the least negative carbon bound to N4 (C7, -0.00445; C5, -0.21717; C21, -0.24162), and the charge on C7–H (+0.30252) is more positive than those on C5 and C21 protons. To the extent that pK_a and natural charge (Q_A, Q_H) are correlated,¹² NPA suggests that C7–H, which is *syn*-coplanar to the N–O_{acyl} bond (11.5°), has the lowest pK_a of the C–H bonds adjacent to N4 (Figures 2 and S1B). Thus, reaction at C7 of IV to form iminium II could occur through an E1-like or E1cb-like transition state by means of the paired TFA.¹³ An alternative to the bimolecular pathway would be the entropically favored *syn* elimination arising from the intramolecular deprotonation of C7–H by the pendant *N*-trifluoroacetyl moiety.





Figure 2. Charges on C and H from the natural population analysis of *N*-acyloxyammonium ion IV.

The mechanistic nuances between the classical Polonovski reaction (acetic anhydride, Ac₂O) and the more reactive Potier variant (TFAA) prompted us to investigate the effect of both acylating agent and temperature on the regioselective functionalization of strychnine *N*-oxide.¹³ The results of those experiments are summarized in Table 1.

Reactions with TFAA (entries 1–3) were less sensitive to temperature such that both percent conversion and ratios of 8 to 7 remained fairly constant. This can be rationalized by considering the nucleofugacity of trifluoroacetate, which greatly favors iminium ion formation.¹³

On the other hand, reactions with Ac₂O (entries 4–6) gave higher ratios of desired carbinolamine 8 to 7. Moreover, they proceeded at a slower rate and required higher temperature for the conversion of starting material to products. The highest ratio



Scheme 3. Syntheses of (–)-sungucine (1), (–)-isosungucine (2), and (–)-strychnogucine B (3).

of **8** to **7** (3.1:1) was obtained at 35 °C for 4 h (entry 6). These results are consistent with an E2 mechanism whereby the more basic acetate can remove the *anti*-periplanar C5–H_β from an *N*-acyloxyammonium strychnine intermediate (Figure 1) to afford the desired carbinolamine **8**.

Table 1. Effect of reagent and temperature on the formation of **7** and **8**.

Entry	Acylating Agent ^[a]	Temperature (°C)	Time (hours) ^[b]	Percent Conversion ^[c]	Ratio of 7 to 8 ^[d]
1	TFAA	0	2	49%	1 : 1.6
2	TFAA	23	2	58%	1 : 1.4
3	TFAA	35	2	52%	1 : 1.9
4	Ac ₂ O	0	4	21%	1 : 1.3
5	Ac ₂ O	23	4	41%	1 : 2.7
6	Ac ₂ O	35	4	62%	1 : 3.1

[a] Two equivalents of the reagent were used using CH₂Cl₂ as solvent.

[b] Iminium ion was trapped with aqueous KOH after 2 or 4 hours.

[c] Percent conversion calculated by ¹H NMR.

[d] Ratio of **7** to **8** determined by ¹H NMR.

With both northern and southern fragments in hand, we focused on forging the characteristic C23–C5' bond in **1–3** using a biomimetic Mannich reaction.¹⁴ After considerable experimentation with various metalation conditions of **5** and screening Lewis acids, we were pleased to find favorable conditions for the Mannich-mediated fragment coupling of **5** and **9**. In the event, **5** was first treated with LDA to generate the lithium enolate (Scheme 3). Addition thereof to the iminium derived from BF₃·Et₂O-treatment of **9** smoothly delivered **10** as a mixture of diastereomers at C23.¹⁵ Subsequent treatment of **10** with HF·Pyridine served to remove the TBS group and effect equilibration at C23 to afford **11** as a single diastereomer; moreover, X-ray analysis of **11** confirmed the stereo- and regiochemical course of the transformations.

The synthesis of (–)-strychnogucine B (**3**) required olefin isomerization from the trisubstituted C16–C17 position into conjugation with the carbonyl. Fortunately, Magnus had reported a similar reaction during early synthetic studies toward strychnine.¹⁶ Thus, treatment of **11** with Cs₂CO₃ in deaerated *tert*-butanol at 85 °C furnished (–)-strychnogucine B (**3**) in 70% yield. Spectral data for **3** (e.g., ¹H and ¹³C NMR, IR) were in complete agreement with those reported in the literature.^{4b}

The syntheses of (–)-sungucine (**1**) and (–)-isosungucine (**2**) from (–)-**3** required two tasks: (1) rupture of the oxepene ring in the southern portion of (–)-strychnogucine B and (2) global deoxygenation of the attendant hydroxyethylidene moieties. The first task was addressed by employing tactics used for northern fragment **5**. Successful execution thereof would deliver two alkene regioisomers (i.e., conjugated and nonconjugated) that map onto **1** and **2**, respectively. In the event, (–)-strychnogucine B (**3**) was first treated with DBU in NMP at 200 °C to afford two bis-diols as a mixture of alkene regioisomers. Subsequent acetylation of this mixture with acetic anhydride to facilitate purification yielded **12** and **13** in a ratio of 1:2 (63–70% overall yield).

Endgame for the syntheses of **1–2** called for adjusting the oxidation states at C18 and C18' in **12–13**. We screened a battery of deoxygenation protocols (e.g., Pd-catalyzed deoxygenation,¹⁷ MsCl/LiEt₃BH,¹⁸ thiocarbonyl diimidazole/Bu₃SnH),¹⁹ but none yielded satisfactory results. Success was ultimately realized by first converting the allylic acetates into their corresponding bromides under acidic conditions and subsequent reduction with NaBH₃CN in DMF.²⁰ Thus, a mixture of **12** and **13** was treated sequentially with 33% HBr followed by NaBH₃CN to furnish (–)-sungucine (**1**) and (–)-isosungucine (**2**) in 40–50% overall yield following chromatographic separation. Spectra data for **1** and **2** (e.g., ¹H and ¹³C NMR, IR) were in complete agreement with the reported values.^{1,4}

In summary, we have completed the first syntheses of (–)-sungucine (**1**), (–)-isosungucine (**2**), and (–)-strychnogucine B (**3**) from commercially available (–)-strychnine (**4**). Key steps

included (1) a DBU-mediated synthesis of isostrychnine from **4**; (2) the Polonovski–Potier reaction of strychnine *N*-oxide; (3) a $\text{BF}_3 \cdot \text{Et}_2\text{O}$ -mediated, biomimetic Mannich coupling; (4) a Cs_2CO_3 -mediated alkene isomerization; and, (5) a NaBH_3CN -mediated reduction of allylic bromides intermediates to install the requisite ethylidene moieties found in targets **1–3**.

Acknowledgements

This research was supported by the NSF (CHE-1111558, CHE-1362461, CNS-09-58854, DEG-12262 awarded to C.N.T.). We thank Prof. Michel Frédérick (Université de Liège) for kindly providing natural samples of (–)-**1-3**. We thank Prof. Scott Bur (Gustavus Adolphus College), Prof. David Dalton (Temple University), and Prof. Jeffrey Winkler (University of Pennsylvania) for helpful discussions. Finally, we kindly dedicate this paper to Prof. Stephen F. Martin on the occasion of his 70th birthday.

Keywords: sungucine • bis-*Strychnos* alkaloids • Polonovski–Potier • biomimetic synthesis • strychnine

- [1] D. J. Newman, G. M. Cragg, *J. Nat. Prod.* **2007**, *70*, 461.
- [2] J. Lamotte, L. Dupont, O. Dideberg, K. Kambu, L. Angenot. *Tetrahedron Lett.* **1979**, *20*, 4227.
- [3] L. Dupont, O. Dideberg, J. Lamotte, K. Kambu, L. Angenot. *Acta Crystallogr. Sect. B-Struct. Sci.* **1980**, *36*, 1669.
- [4] a) M. Frederich, M.-C. De Pauw, G. Llabres, M. Tits, V. Brandt, J. Penelle, P. De Mol, L. Angenot. *Planta. Med.* **2000**, *66*, 262; b) M. Frederich, M.-C. De Pauw, C. Prosperi, M. Tits, V. Brandt, J. Penelle, M.-P. Hayette, P. De Mol, L. Angenot. *J. Nat. Prod.* **2001**, *64*, 12.
- [5] A. Lansiaux, C. Bailly, C. Houssier, P. Colson, M. C. De Pauw-Gillet, M. Frederich, M. Tits, L. Angenot. *Planta. Med.* **2002**, *68*, 591.
- [6] a) M. Frederich, M. Bentires-Ali, M. Tits, L. Angenot, E. Heinen, M. C. De Pauw-Gillet. *Naunyn-Schmiedebergs Arch. Pharmacol.* **2003**, *367*, 260; b) M. Frederich, M. Tits, L. Angenot. *Chem. Nat. Compd.* **2003**, *39*, 513.
- [7] M. Frederich, M. J. Jacquier, P. Thepenier, P. De Mol, M. Tits, G. Philippe, C. Delaude, L. Angenot, M. Zeches-Hanrot. *J. Nat. Prod.* **2002**, *65*, 1381.
- [8] G. F. Smith in *The Alkaloids: Chemistry and Physiology*, Vol. 8 (Ed: R. H. F. Manske), Academic Press, **1965**, pp. 591–671.
- [9] A. Ahond, A. Cave, C. Kan-Fan, P. H. Husson, J. De Rostolan, P. Potier. *J. Am. Chem. Soc.* **1968**, *90*, 5622.
- [10] E. Clot. *Eur. J. Inorg. Chem.* **2009**, *2009*, 2319.
- [11] A. E. Reed, R. B. Weinstock, F. Weinhold. *J. Chem. Phys.*, **1985**, *83*, 735.
- [12] P. G. Seybold in *Advances in Quantum Chemistry*, vol. 64 (Eds: R. S. John, J. B. Erkki), Academic Press, **2012**, pp. 83–104.
- [13] S. K. Bur in *Comprehensive Organic Synthesis II (Second Edition)* (Ed: P. Knochel), Elsevier, **2014**, pp. 755–801.
- [14] M. Arend, B. Westermann, N. Risch, *Angew. Chem. Int. Ed.* **1998**, *37*, 1044; *Angew. Chem.* **1998**, *110*, 1096.
- [15] Attempts to determine the exact diastereomeric ratio by ¹H NMR was unsuccessful due to the configurational lability at C23.
- [16] P. Magnus, M. Giles, R. Bonnert, G. Johnson, L. McQuire, M. Deluca, A. Merritt, C. S. Kim, N. Vicker, *J. Am. Chem. Soc.* **1993**, *115*, 8116.
- [17] J. Tsuji, T. Mandai, *Synthesis*, **1996**, 1996, 1.
- [18] N. Kogure, Y. Suzuki, Y. Wu, M. Kitajima, R. Zhang, H. Takayama, *Tetrahedron Lett.* **2012**, *53*, 6523.
- [19] K. C. Nicolaou, C. K. Hwang, A. L. Smith, S. V. Wendebom, *J. Am. Chem. Soc.* **1990**, *112*, 7416.
- [20] L. Fernandes, A. J. Bortoluzzi, M. M. Sa, *Tetrahedron* **2004**, *60*, 9983.

Entry for the Table of Contents (Please choose one layout)

Layout 1:

COMMUNICATION

Text for Table of Contents

((Insert TOC Graphic here))

Author(s), Corresponding Author(s)*

Page No. – Page No.

Title

Layout 2:


COMMUNICATION

*Senzhi Zhao, Christiana N. Tejjaro,
Heng Chen, Gopal Sirasani, Shivaiah
Vaddypally, Michael J. Zdilla, Graham E.
Dobereiner, and Rodrigo B. Andrade**

Page No. – Page No.

**Concise Syntheses of Bis-Strychnos
Alkaloids (–)-Sungucine, (–)-
Isosungucine, and (–)-Strychnogucine B
from (–)-Strychnine**

Text for table of contents



[a] S. Zhao, C. N. Tejjaro, H. Chen, G. Sirasani, S. Vaddypally, Zdilla, G. E. Dobereiner, R. B. Andrade
Department of Chemistry
Temple University
1901 N. 13th St. Philadelphia, Pa 19122
E-mail: randrade@temple.edu

Supporting information for this article is given via a link at the end of the document.