



Some characteristics of hyperglycaemic crisis differ between patients with and without COVID-19 at a safety-net hospital in a cross-sectional study

Arnav Shah, Andrew Deak, Shaneisha Allen, Elayna Silfani, Christina Koppin, Yaara Zisman-Ilani, Imali Sirisena, Christina Rose & Daniel Rubin

To cite this article: Arnav Shah, Andrew Deak, Shaneisha Allen, Elayna Silfani, Christina Koppin, Yaara Zisman-Ilani, Imali Sirisena, Christina Rose & Daniel Rubin (2021) Some characteristics of hyperglycaemic crisis differ between patients with and without COVID-19 at a safety-net hospital in a cross-sectional study, *Annals of Medicine*, 53:1, 1642-1645, DOI: [10.1080/07853890.2021.1975042](https://doi.org/10.1080/07853890.2021.1975042)

To link to this article: <https://doi.org/10.1080/07853890.2021.1975042>



© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 11 Sep 2021.



[Submit your article to this journal](#)



Article views: 467



[View related articles](#)



[View Crossmark data](#)

CLINICAL STUDY



Some characteristics of hyperglycaemic crisis differ between patients with and without COVID-19 at a safety-net hospital in a cross-sectional study

Arnav Shah^a, Andrew Deak^a, Shaneisha Allen^b, Elayna Silfani^c, Christina Koppin^a, Yaara Zisman-Ilani^d, Imali Sirisena^b, Christina Rose^c and Daniel Rubin^b

^aLewis Katz School of Medicine, Temple University, Philadelphia, PA, USA; ^bSection of Endocrinology, Diabetes, and Metabolism, Lewis Katz School of Medicine at Temple University, Philadelphia, PA, USA; ^cSchool of Pharmacy, Temple University, Philadelphia, PA, USA; ^dDepartment of Social and Behavioral Sciences, College of Public Health, Temple University, Philadelphia, PA, USA

ABSTRACT

Objective: To compare patients with DKA, hyperglycaemic hyperosmolar syndrome (HHS), or mixed DKA-HHS and COVID-19 [COVID (+)] to COVID-19-negative (–) [COVID (–)] patients with DKA/HHS from a low-income, racially/ethnically diverse catchment area.

Methods: A cross-sectional study was conducted with patients admitted to an urban academic medical center between 1 March and 30 July 2020. Eligible patients met lab criteria for either DKA or HHS. Mixed DKA-HHS was defined as meeting all criteria for either DKA or HHS with at least 1 criterion for the other diagnosis.

Results: A total of 82 participants were stratified by COVID-19 status and type of hyperglycaemic crisis [26 COVID (+) and 56 COVID (–)]. A majority were either Black or Hispanic. Compared with COVID (–) patients, COVID (+) patients were older, more Hispanic and more likely to have type 2 diabetes (T2D, 73% vs 48%, $p < .01$). COVID(+) patients had a higher mean pH (7.25 ± 0.10 vs 7.16 ± 0.16 , $p < .01$) and lower anion gap (18.7 ± 5.7 vs 22.7 ± 6.9 , $p = .01$) than COVID (–) patients. COVID (+) patients were given less intravenous fluids in the first 24 h (2.8 ± 1.9 vs 4.2 ± 2.4 L, $p = .01$) and were more likely to receive glucocorticoids (95% vs. 11%, $p < .01$). COVID (+) patients may have taken longer to resolve their hyperglycaemic crisis (53.3 ± 64.8 vs 28.8 ± 27.5 h, $p = .09$) and may have experienced more hypoglycaemia < 3.9 mmol/L (35% vs 19%, $p = .09$). COVID (+) patients had a higher length of hospital stay (LOS, 14.8 ± 14.9 vs 6.5 ± 6.0 days, $p = .01$) and in-hospital mortality (27% vs 7%, $p = .02$).

Discussion: Compared with COVID (–) patients, COVID (+) patients with DKA/HHS are more likely to have T2D. Despite less severe metabolic acidosis, COVID (+) patients may require more time to resolve the hyperglycaemic crisis and experience more hypoglycaemia while suffering greater LOS and risk of mortality. Larger studies are needed to examine whether differences in management between COVID (+) and (–) patients affect outcomes with DKA/HHS.

ARTICLE HISTORY

Received 28 May 2021
Accepted 26 August 2021

KEYWORDS


COVID-19; hyperglycaemic emergencies; diabetic ketoacidosis; hyperglycaemic hyperosmolar syndrome

Introduction

The clinical presentation and outcomes of patients with COVID-19 and hyperglycaemic emergency (i.e. diabetic ketoacidosis [DKA] or hyperglycaemic hyperosmolar syndrome [HHS]) have been described [1–3]. Few studies have directly compared patients in DKA with COVID-19 to those without COVID-19 [4–6] and none have specifically studied patients at a safety-net hospital. The objective of this study was to compare the clinical characteristics, management, and outcomes of patients with DKA, HHS, or mixed DKA-HHS and a positive SARS-CoV-2 RT-PCR to negative patients with DKA/HHS from a low-income, racially and ethnically diverse catchment area.

Patients and Methods

A cross-sectional study was conducted with patients admitted to an urban academic medical Centre in Philadelphia, PA, USA between 1 March and 30 July 2020. Data were collected by computer query and manual review of electronic health records. Selection bias was limited by enrolling all eligible patients. Patients met lab criteria for either DKA (serum bicarbonate ≤ 18 mEq/L, anion gap > 10 , ketonaemia or ketonuria, and arterial pH ≤ 7.30), or HHS (blood glucose > 33.3 mmol/L and osmolality > 320 mOsm/kg) [7]. Mixed DKA-HHS was defined as meeting all criteria for either DKA or HHS with at least 1 criterion for the other diagnosis. Diagnoses were adjudicated by an

CONTACT Daniel Rubin  daniel.rubin@tuhs.temple.edu  Lewis Katz School of Medicine at Temple University, Section of Endocrinology, Diabetes, and Metabolism, Philadelphia, PA, USA

This article has been republished with minor changes. These changes do not impact the academic content of the article.

© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 1. Continued.

Variable	All Cases ^a	COVID Positive				COVID Negative				p Value ^c
		All ^a	DKA ^b	HHS ^b	Mixed DKA/HHS ^b	All ^a	DKA ^b	HHS ^b	Mixed DKA/HHS ^b	
<3.9 mmol/L (70 mg/dl)	19 (23%)	9 (35%)	7 (35%)	1 (33%)	1 (33%)	10 (18%)	7 (19%)	1 (10%)	2 (20%)	.09
<3.0 mmol/L (54 mg/dl)	5 (6.1%)	2 (8%)	1 (5%)	0 (0%)	1 (33%)	3 (5%)	1 (2.8%)	1 (10%)	1 (10%)	.68
<2.2 mmol/L (40 mg/dl)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	–
Outcomes										
Length of hospital stay, d	9.1 ± 10.4	14.8 ± 14.9	10.6 [7.0–19.3]	5.9 [5.3–13.3]	6.8 [5.2–9.4]	6.5 ± 6.0	4.2 [2.1–8.2]	5.6 [3.1–9.8]	5.3 [4.3–8.9]	.01
ICU stay, n (%)	58 (72%)	17 (65%)	12 (60%)	2 (67%)	3 (100%)	41 (75%)	26 (72%)	7 (70%)	8 (89%)	.39
In-hospital mortality, n (%)	11 (14%)	7 (27%)	6 (30%)	1 (33%)	0 (0%)	4 (7%)	2 (6%)	1 (10%)	1 (10%)	.02

^aData are n (%) or mean ± SD; ^bData are n (%) or median [IQR]; ^cAll COVID positive vs. All COVID negative; “–” indicates data not available.

endocrinologist. Continuous variables were compared between COVID-19 positive patients [at least one positive SARS-CoV-2 RT-PCR, COVID (+)] and negative patients [all negative SARS-CoV-2 RT-PCR or not tested, COVID (–)] with *t*-tests and categorical variables were compared with chi-square. For DKA/HHS subgroup comparisons across COVID-19 status, ANOVA, chi-square, or Fisher Exact tests were used. The Temple University Institutional Review Board approved the study (number 27246). The requirement for informed consent was waived because this was a study of pre-existing retrospective data.

Results

A total of 82 participants were eligible and stratified by COVID-19 status and type of hyperglycaemic crisis (26 COVID (+) and 56 COVID (–), Table 1). A majority of the patients were either Black or Hispanic. The distribution of hyperglycaemic emergency type was not significantly different between COVID (+) and COVID (–) groups. Compared with COVID (–) patients, COVID (+) patients were older (mean age 58.0 ± 14.3 vs 50.0 ± 18.0 years, *p* = .05), identified more as Hispanic (35% vs 15%, *p* < .05) and were more likely to have type 2 diabetes (T2D, 73% vs 48%, *p* < .01). Among those with DKA, T2D was more common in COVID (+) than in COVID (–) patients, whereas T1D was more common in COVID (–) patients (*p* = .02). Additionally, COVID (+) DKA patients had less severe ketoacidosis with a higher pH (*p* = .03) and bicarbonate (*p* < .01) and lower anion gap (*p* < .01) than COVID (–) patients. Overall, COVID (+) patients had a higher mean pH (7.25 ± 0.10 vs 7.16 ± 0.16, *p* < .01) and lower anion gap (18.7 ± 5.7 vs 22.7 ± 6.9, *p* = .01) than COVID (–) patients. COVID (+) patients were given less intravenous fluids in the first 24 h after admission (2.8 ± 1.9 vs 4.2 ± 2.4 L, *p* = .01) and were more likely to receive glucocorticoids (95% vs. 11%, *p* < .01). COVID (+) patients may have taken longer to resolve their hyperglycaemic crisis (53.3 ± 64.8 vs 28.8 ± 27.5 h, *p* = .09) and may have experienced more hypoglycaemia <3.9 mmol/L (35% vs 19%, *p* = .09). COVID (+) patients had a higher length of hospital stay (LOS, 14.8 ± 14.9 vs 6.5 ± 6.0 days, *p* = .01) and in-hospital mortality (27% vs 7%, *p* = .02).

Discussion

Compared with COVID (–) patients, COVID (+) patients with DKA/HHS are more likely to have T2D. Despite less severe metabolic acidosis, COVID (+) patients may require more time to resolve the hyperglycaemic crisis and experience more hypoglycaemia while requiring

longer hospital stays and suffering a greater risk of mortality. Whether or not the difference in clinical management in terms of intravenous fluids and glucocorticoids is associated with outcomes remains unclear.

While other studies have described the association of COVID-19 with T2D, higher LOS, and greater mortality among DKA patients,^[4–6] metabolic differences from COVID (-) patients have not been found consistently. We are unaware of other studies comparing COVID (+) and COVID (-) patients on such an extensive list of variables across types of hyperglycaemic emergencies. Furthermore, this study uniquely focussed on racially and ethnically diverse patients at a safety-net hospital. These findings may not be generalizable to other populations, and the study is limited by a modest sample size drawn from a single center. Larger studies are needed to examine whether differences in management between COVID (+) and COVID (-) patients affect outcomes with DKA/HHS.

Acknowledgments

This study was not funded. All authors had full access to all the data in the study and accept responsibility to submit for publication. Publication of this article was funded in part by the Temple University Libraries Open Access Publishing Fund.

Disclosure statement

DR has received research funding from AstraZeneca and NIH/NIDDK (R01DK122073) for unrelated work. The other authors report no conflict of interest.

Data availability statement

The data that support the findings of this study are available from the corresponding author, DR, upon reasonable request.

References

- [1] Armeni E, Aziz U, Qamar S, et al. Protracted keto-naemia in hyperglycaemic emergencies in COVID-19: a retrospective case series. *Lancet Diabetes Endocrinol.* 2020;8(8):660–663.
- [2] Papadopoulou VP, Koutroulos MV, Zikoudi DG, et al. Diabetes-related acute metabolic emergencies in COVID-19 patients: a systematic review and Meta-analysis. *Diabetol Int.* 2021;23:1–15.
- [3] Pal R, Banerjee M, Yadav U, et al. Clinical profile and outcomes in COVID-19 patients with diabetic ketoacidosis: a systematic review of literature. *Diabetes Metab Syndr.* 2020;14(6):1563–1569.
- [4] Pasquel FJ, Messler J, Booth R, et al. Characteristics of and mortality associated with diabetic ketoacidosis among US patients hospitalized with or without COVID-19. *JAMA Netw Open.* 2021;4(3):e211091.
- [5] Kempegowda P, Melson E, Johnson A, et al. Effect of COVID-19 on the clinical course of diabetic ketoacidosis (DKA) in people with type 1 and type 2 diabetes. *Endocr Connect.* 2021;10(4):371–377.
- [6] Misra S, Khozoe B, Huang J, et al. Comparison of diabetic ketoacidosis in adults during the SARS-CoV-2 outbreak and over the same time period for the preceding 3 years. *Diabetes Care.* 2021;44(2):e29–e31.
- [7] Kitabchi AE, Umpierrez GE, Miles JM, et al. Hyperglycemic crises in adult patients with diabetes. *Diabetes Care.* 2009;32(7):1335–1343.